



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/32180 (43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/US99/28483 (22) International Filing Date: 1 December 1999 (01.12.99) (30) Priority Data: 60/110,395 1 December 1998 (01.12.98) US (71) Applicant (for all designated States except US): THE INSTITUTES FOR PHARMACEUTICAL DISCOVERY, INC. [US/US]; 23 Business Park Drive, Branford, CT 06405 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SREDY, Janet [US/US]; 98 Trumbull Avenue, Milford, CN 06460 (US). JACOT, Jorge [US/US]; 2 Katie-Joe Lane, Branford, CN 06405 (US). (74) Agent: SARUSSI, Steven, J.; McDonnell, Boehnen, Hulbert & Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHODS OF REDUCING SERUM GLUCOSE AND TRIGLYCERIDE LEVELS AND FOR INHIBITING ANGIOGENESIS USING SUBSTITUTED INDOLEALKANOIC ACIDS (57) Abstract Disclosed are methods of reducing serum glucose and triglyceride levels and for inhibiting angiogenesis, the methods comprising administration of substituted indolealkanoic acids to patients in need of such treatment. Also disclosed are such compounds useful in the treatment of angiogenesis, hyperglycemia, hyperlipidemia and chronic complications arising from diabetes mellitus. Also disclosed are pharmaceutical compositions containing the compounds.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS OF REDUCING SERUM GLUCOSE AND
TRIGLYCERIDE LEVELS AND FOR INHIBITING ANGIOGENESIS
USING SUBSTITUTED INDOLEALKANOIC ACIDS

5

Background of Invention:

The use of aldose reductase inhibitors (ARIs) for the treatment of diabetic complications is well known. The complications arise from elevated levels of glucose in tissues
10 such as the nerve, kidney, retina and lens that enters the polyol pathway and is converted to sorbitol via aldose reductase. Because sorbitol does not easily cross cell membranes, it accumulates inside certain cells resulting in changes in osmotic pressure, alterations in the redox state of
15 pyridine nucleotides (i.e. increased NADH/NAD⁺ ratio) and depleted intracellular levels of myoinositol. These biochemical changes, which have been linked to diabetic complications, can be controlled by inhibitors of aldose reductase.

20 The use of aldose reductase inhibitors for the treatment of diabetic complications has been extensively reviewed, see:
(a) *Textbook of Diabetes*, 2nd ed.; Pickup, J. C. and Williams, G. (Eds.); Blackwell Science, Boston, MA 1997.; (b) Larson, E. R.; Lipinski, C. A. and Sarges, R., *Medicinal Research Reviews*,
25 1988, 8 (2), 159-198; (c) Dvornik, D. *Aldose Reductase Inhibition*. Porte, D. (ed), Biomedical Information Corp., New York, NY. Mc Graw Hill 1987; (d) Petrash, J. M., Tarle, I.,

Wilson, D. K. Quioco. F. A. *Perspectives in Diabetes, Aldose Reductase Catalysis and Crystallography: Insights From Recent Advances in Enzyme Structure and Function*, Diabetes, 1994, 43, 955; (e) Aotsuka, T.; Abe, N.; Fukushima, K.; Ashizawa, N. and
5 Yoshida, M., *Bioorg. & Med. Chem. Letters*, 1997, 7, 1677, (f)
, T., Nagaki, Y.; Ishii, A.; Konishi, Y.; Yago, H; Seishi, S.;
Okukado, N.; Okamoto, K., *J. Med. Chem.*, 1997, 40, 684; (g)
Ashizawa, N.; Yoshida, M.; Sugiyama, Y.; Akaike, N.; Ohbayashi,
S.; Aotsuka, T.; Abe, N.; Fukushima, K.; Matsuura, A, *Jpn. J.*
10 *Pharmacol.* 1997, 73, 133; (h) Kador, P. F.; Sharpless, N. E.,
Molecular Pharmacology, 1983, 24, 521; (I) Kador, P. F.;
Kinoshita, J. H.; Sharpless, N. E., *J. Med. Chem.* 1985, 28 (7),
841; (j) Hotta, N., *Biomed. & Pharmacother.* 1995, 5, 232; (k)
Mylar, B.; Larson, E. R.; Beyer, T. A.; Zembrowski, W. J.;
15 Aldinger, C. E.; Dee, F. D.; Siegel, T. W.; Singleton, D. H.,
J. Med. Chem. 1991, 34, 108; (l) Dvornik, D. *Croatica Chemica*
Acta 1996, 69 (2), 613.

Previously described aldose reductase inhibitors most
closely related to the present invention include those sighted
20 in: (a) U.S Pat. No. 5,700,819: 2-Substituted benzothiazole
derivatives useful in the treatment of diabetic complications,
(b) U.S Pat. No. 4,868,301: Processes and intermediates for the
preparation of oxophthalazinyll acetic acids having
benzothiazole or other heterocyclic side chains, (c) U.S Pat.
25 No. 5,330,997: 1H-indazole-3-acetic acids as aldose reductase

inhibitors, and (d) U.S. Pat. No. 5,236,945: 1H-indazole-3-acetic acids as aldose reductase inhibitors. Although many aldose reductase inhibitors have been extensively developed, none have demonstrated sufficient efficacy in human clinical trials without significant undesirable side effects. Thus no aldose reductase inhibitors are currently available as approved therapeutic agents in the United States; and consequently, there is still a significant need for new, efficacious and safe medications for the treatment of diabetic complications.


Treatment to normalize the plasma glucose concentration in people afflicted with type 2 diabetes currently includes diet, exercise and oral agents such as sulfonylureas, metformin and glitazone-type compounds. Many of these agents exhibit side effects and have limited efficacy. There is a need for new agents which do not possess these drawbacks. Because of the limited efficacy of each method of treatment often the oral agents are giving in combination of with each other or with insulin.

Elevated serum triglyceride levels are also commonly associated with diabetes; however, this condition is also widely seen in nondiabetic patients. The mechanism causing the presence of elevated triglyceride levels in patients, both diabetic and otherwise, is different from that underlying chronic diabetes-related complications directly treatable by inhibition of aldose reductase activity. There is, therefore,

a need for treatment of elevated triglyceride levels in diabetic and/or nondiabetic patients, e.g., cardiac patients.

Summary of the Invention:

This invention provides compounds that interact with and inhibit aldose reductase. Thus, in a broad aspect, the invention provides compounds of Formula I:



I

or pharmaceutically acceptable salts thereof wherein

A is a C₁-C₄ alkylene group optionally substituted with C₁-C₂ alkyl or mono- or disubstituted with halogen, preferably fluoro or chloro;

Z is a bond, O, S, C(O)NH, or C₁-C₃ alkylene optionally substituted with C₁-C₂ alkyl;

R₁ is hydrogen, alkyl having 1-6 carbon atoms, halogen, 2-, 3-, or 4-pyridyl, or phenyl, where the phenyl or pyridyl is optionally substituted with up to three groups selected from halogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl, nitro, amino, or mono- or di(C₁-C₆)alkylamino;

R₂, R₃, R₄ and R₅ are each independently

hydrogen, halogen, nitro, or an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens);

OR₇, SR₇, S(O)R₇, S(O)₂(R₇)₂, C(O)N(R₇)₂, or N(R₇)₂, wherein

each R_i is independently hydrogen, an alkyl group of 1-

6 carbon atoms (which may be substituted with one or

more halogens) or benzyl, where the phenyl portion is

optionally substituted with up to three groups

independently selected from halogen, C₁-C₆ alkyl, C₁-C₆

alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenyl

or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-,

3-, or 4-pyridyl, each of which phenyl or heteroaryl

is optionally substituted with up to three groups

independently selected from halogen, C₁-C₆ alkyl, C₁-C₆

alkoxy, amino, and mono- or di(C₁-C₆)alkylamino;


phenoxy where the phenyl portion is optionally substituted

with up to three groups independently selected from

halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or

di (C₁-C₆) alkylamino; or

a group of the formula



where

J is a bond, CH₂, oxygen, or nitrogen; and

each r is independently 2 or 3;

R₆ is hydroxy or a prodrug group;

R_a is hydrogen, C₁-C₆ alkyl, fluoro, or trifluoromethyl; and

Ar represents aryl or heteroaryl, each of which is optionally substituted with up to five groups.

In another aspect, the invention provides methods for
5 preparing such compounds.

The compounds of the invention inhibit aldose reductase. Since aldose reductase is critical to the production of high levels of sorbitol in individuals with diabetes, inhibitors of aldose reductase are useful in preventing and/or treating
10 various complications associated with diabetes. The compounds of the invention are therefore effective for the treatment of diabetic complications as a result of their ability to inhibit aldose reductase.

In another aspect, the invention provides methods for
15 treating and/or preventing chronic complications associated with diabetes mellitus, including, for example, diabetic cataracts, retinopathy, keratopathy, wound healing, diabetic uveitis, diabetic cardiomyopathy, nephropathy, and neuropathy.

The compounds of this invention also possess
20 antihyperglycemic activity and are therefore useful for the treatment of hyperglycemia. and elevated serum triglyceride levels. Accordingly, an aspect of the invention is prevention and/or alleviation of complications associated with hyperglycemia with the inventive compounds.

The compounds of the present invention have been discovered to lower triglycerides. While serum triglyceride levels are often elevated in diabetic patients, they are also frequently elevated in nondiabetic patients resulting in various diseases and disorders, e.g., cardiac disease. Because of their ability to reduce serum triglyceride levels, the compounds of the present invention are useful in the treatment, i.e., prevention and/or alleviation, of elevated triglyceride levels in both diabetic and nondiabetic patients.

Thus, the compounds of the present invention may be used as antihyperlipidemic and/or antihyperglycemic agents. The compounds of this invention may be given in combination with other glucose or lipid lowering agents as well as other agents that are given specifically to treat the complications of diabetes.

It has also been discovered that the compounds of the present invention exhibit anti-angiogenic activity in an established in vitro assay. The discovery of this biological activity for the compounds of the invention is unexpected. As a result of this biological activity, the compounds of the invention can be used to treat various diseases that exhibit aberrant vasoproliferation. According to the invention, the compound would be administered to a mammal in need of inhibition of vasoproliferation, i.e., inhibition of angiogenesis. Examples of such diseases are diabetic

retinopathy, age-related macular degeneration, retinopathy of prematurity, corneal neovascularization, pterygium, and any neoplasms (cancers) which appear to be angiogenesis dependent. Administration of the compound(s) of this invention is/are not
5 limited to a particular mode, and could be administered systemically or topically to the eye in an appropriate ophthalmic solution. The compounds of the invention may be administered in combination therapy with other known anti-angiogenic agents.

10 The compounds of the invention have also been discovered to promote the healing of wounds in mammals. In preferred aspects, the compounds are useful in promoting wound healing in diabetic mammals. Thus, the compounds of the invention may be employed in the treatment of wounds in mammals, preferably
15 humans, more preferably in diabetic humans.


In still another aspect, the invention provides pharmaceutical compositions containing compounds of Formula I.

In still another aspect, the invention provides for the use of a compound or compounds of Formula I for the preparation
20 of a medicament for the treatment of any of the disorders or diseases (a) listed above, (b) connected with diabetic complications, hyperglycemia, or hypertriglyceridemia, or (c) where inhibition of vasoproliferation is indicated.

Detailed Description of the Invention

As used herein, the term "treatment" includes both prevention and alleviation.

The numbering system for the compounds of Formula I is as follows:



I

As noted above, the invention provides novel substituted indole alkanolic acids useful in treating and/or preventing complications associated with or arising from elevated levels of glucose in individuals suffering from diabetes mellitus. These compounds are represented by Formula I above.

In compounds of Formula I, the aryl and heteroaryl groups represented by Ar include:

a phenyl group optionally substituted with up to 5 groups independently selected from halogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens), nitro, OR₇, SR₇, S(O)R₇, S(O)₂R₇, or N(R₇)₂, wherein R₇ is hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆

alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino, or the phenyl group may be condensed with benzo where the benzo is optionally substituted with one or two of halogen, cyano, nitro, trifluoromethyl, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl;

a heterocyclic 5-membered ring having one nitrogen, oxygen or sulfur, two nitrogens one of which may be replaced by oxygen or sulfur, or three nitrogens one of which may be replaced by oxygen or sulfur, said heterocyclic 5-membered ring substituted by one or two fluoro, chloro, (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo, cyano, nitro, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, one or two of fluoro, chloro, bromo, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl or trifluoromethyl, or two fluoro or two trifluoromethyl with one hydroxy or one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one trifluoromethyl, or three fluoro, said pyridyl, furyl

or thienyl optionally substituted in the 3-position by fluoro, chloro, bromo, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

a heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur, said heterocyclic 6-membered ring substituted by one or two (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo or trifluoromethylthio, or one or two of fluoro, chloro, bromo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, or trifluoromethyl, and said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

said benzo-condensed heterocyclic 5-membered or 6-membered rings optionally substituted in the heterocyclic 5-membered or 6-membered ring by one of fluoro, chloro, bromo, methoxy, or trifluoromethyl;

oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms, with thiophene or with furane, each optionally substituted by one of fluoro, chloro, bromo, trifluoromethyl, methylthio or methylsulfinyl;

imidazolopyridine or triazolopyridine optionally substituted by
one of trifluoromethyl, trifluoromethylthio, bromo, or
(C₁-C₆)alkoxy, or two of fluoro or chloro;
thienothiophene or thienofuran optionally substituted by one of
5 fluoro, chloro or trifluoromethyl; thienotriazole
optionally substituted by one of chloro or
trifluoromethyl;
naphthothiazole; naphthoxazole; or thienoisothiazole.

10 More specific compounds of the invention are those of
Formula I wherein Ar is optionally substituted benzothiazolyl,
benzoxazolyl, isoquinolyl, benzothiophen-yl, benzofuran-yl or
benzimidazolyl, or substituted oxadiazolyl or indolyl. Other
more specific compounds are of Formula I those wherein R_a is
15 trifluoromethyl, Z is a covalent bond or CH₂, R₆ is hydroxy,
and each of R₂-R₅ are independently hydrogen, halogen, more
preferably bromo or chloro, C₁-C₂ alkyl, phenoxy, benzyloxy, or
C₁-C₂ alkoxy, and R₁ is hydrogen or methyl.

Preferred compounds of the invention are those wherein Z
20 is a covalent bond, R₆ is hydroxy, Ar is optionally substituted
benzothiazol-2-yl, benzothiazol-5-yl, benzoisothiazol-3-yl,
benzoxazol-2-yl, 2-quinolyl, 2-quinoxalyl, oxazolo[4,5-
b]pyridine-2-yl, benzothiophen-2-yl, benzofuran-2-yl, or
thiazolo[4,5-pyridine-2-yl, thieno[2,3-b]pyridine-2-yl,
25 imidazo[1,5-a]pyridine-2-yl, or indol-2-yl, or substituted

isothiazol-4-yl, 1,3,4-oxadiazol-5-yl, 1,2,5-thiadiazol-3-yl, oxazol-2-yl, thiazol-2-yl, or thiazol-4-yl, R_2 - R_5 are independently hydrogen, halogen, more preferably bromo or chloro, C_1 - C_2 alkyl, phenoxy, benzyloxy or phenyl where each phenyl portion is optionally substituted with C_1 - C_6 alkyl, halogen, C_1 - C_6 alkoxy, hydroxy, amino or mono- or di (C_1 - C_6) alkylamino R_6 is hydrogen, fluoro or C_1 - C_2 alkyl, and R_1 is hydrogen or methyl.

10 Other preferred compounds are those wherein the methylene bridge connecting the indolyl group with Ar is located alpha with respect to a nitrogen atom in Ar, e.g. wherein Ar is benzoxazol-2-yl or 1,2,4-oxadiazol-3-yl mentioned above.


Other more specific compounds of the invention are those
15 wherein Z is a covalent bond, R_6 is hydroxy, R_7 is hydrogen, Ar is optionally 4,5,6 or 7 benzo-substituted benzothiazolyl, benzoxazolyl, benzimidazolyl, benzothiophenyl, benzofuranyl, or indolyl, or Ar is 2-benzothiazolyl substituted on benzo by one trifluoroacetyl or trifluoromethylthio, or one or two of fluoro
20 chloro, bromo, hydroxy, methyl, methoxy, trifluoromethyl, trifluoromethoxy, trifluoromethylthio, or one or, preferably, two fluoro and one trifluoromethyl, or two fluoro or two trifluoromethyl with one methoxy, or three fluoro, or by 6,7-benzo, and those wherein one of R_2 and R_3 is hydrogen, fluoro,
25 chloro, bromo or methyl, and one of R_4 and R_5 is hydrogen, or

chloro, bromo, methyl, isopropyl, methoxy, nitro or
trifluoromethyl; or R_3 and R_4 is 5, 6-difluoro, R_5 is hydrogen;
and those wherein Ar is optionally substituted benzothiazol-2-
yl or quinoxalyl and R_3 and R_4 are each chloro, and R_1 is
5 hydrogen or methyl.

Further more specific compounds are those wherein Z is a
covalent bond, R_6 is hydroxy, Ar is optionally substituted
benzothiazol-2-yl, R_3 and R_4 are hydrogen, and R_5 is methyl;
those wherein Z is a covalent bond, R_6 is hydroxy, R_3 , R_4 and R_5
10 are hydrogen, chloro, fluoro, bromo or C_1 - C_2 alkyl, R_6 is
hydrogen, and Ar is optionally 4,5,6 or 7 benzosubstituted
benzothiazolyl-2-trifluoromethyl, benzoxazolyl-2-
trifluoromethyl, benzimidazolyl-2-trifluoromethyl, benzofuran-
2-trifluoromethyl, benzofuran-3-trifluoromethyl, benzothiophen-
15 2-trifluoromethyl, benzothiophen-3-trifluoromethyl, indolyl-2-
trifluoromethyl, or indolyl-3-trifluoromethyl; and those
wherein Z is CH_2 , R_6 is hydroxy, Ar is optionally substituted
benzothiazol-2-yl, benzothiazol-5-yl, benzoisothiazol-3-yl,
benzoxazol-2-yl, 2-quinolyl, 2-quinoxalyl, oxazolo[4,5-
20 b]pyridine-2-yl, or thiazolo[4,5-b]pyridine-2-yl, or
substituted 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl,
isothiazol-5-yl, isothiazol-4-yl, 1,3,4-oxadiazol-5-yl, 1,2,5-
thiadiazol-3-yl, oxazol-2-yl, thiazol-2-yl, or thiazol-4-yl,
and R_3 , R_4 and R_5 are independently hydrogen, chloro, fluoro,
25 bromo, C_1 - C_2 alkyl, or trifluoromethyl, and R_6 is hydrogen.

Generally, R_1 in the specific compounds described above is hydrogen, halogen, preferably chloro or fluoro, C_1 - C_6 alkyl, or phenyl optionally substituted with up to three groups independently selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, amino, and mono- or di(C_1 - C_6)alkylamino. Preferred R_1 groups are hydrogen and methyl.

Preferred compounds of the invention include those where Ar in Formula I is substituted phenyl, i.e., compounds of Formula II:



II

wherein

A is a C_1 - C_4 alkylene group optionally substituted with C_1 - C_2 alkyl;

Z is a bond, or C_1 - C_3 alkylene optionally substituted with C_1 - C_2 alkyl;

R_a is hydrogen, C_1 - C_6 alkyl, chloro, bromo, fluoro, or trifluoromethyl;

R_1 is hydrogen, C_1 - C_6 alkyl, fluoro, or phenyl optionally substituted with up to three groups independently selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, amino, and mono- or di(C_1 - C_6)alkylamino;

R_2 , R_3 , R_4 and R_5 are each independently

hydrogen, halogen, an alkyl group of 1-6 carbon atoms

(which may be substituted with one or more halogens),

nitro, OR_7 , SR_7 , $S(O)R_7$, $S(O)_2N(R_7)_2$, $C(O)N(R_7)_2$, or

5 $N(R_7)_2$, wherein each R_7 is independently hydrogen, an

alkyl group of 1-6 carbon atoms (which may be

substituted with one or more halogens) or benzyl,

where the phenyl portion is optionally substituted

with up to three groups independently selected from

10 halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, amino, and mono- or

di(C_1 - C_6)alkylamino;

phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-,

3-, or 4-pyridyl, each of which phenyl or heteroaryl

is optionally substituted with up to three groups

15 independently selected from halogen, C_1 - C_6 alkyl, C_1 - C_6

alkoxy, amino, and mono- or di(C_1 - C_6)alkylamino;


phenoxy where the phenyl portion is optionally substituted

with up to three groups independently selected from

halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, amino, and mono- or

20 di(C_1 - C_6)alkylamino; or

a group of the formula



where

J is a bond, CH_2 , oxygen, or nitrogen; and

each r is independently 2, or 3;


R_6 is hydrogen, an alkoxy group of 1-6 carbon atoms, or $-O^+M^+$

where M^+ is a cation forming a pharmaceutically acceptable salt; and

5 R_8 , R_9 , and R_{10} are independently hydrogen, fluorine, chlorine, bromine, trifluoromethyl or nitro.

Other preferred compounds of the invention are those where Ar is a substituted benzothiazole, i.e., compounds of Formula

10 III:



III

wherein

A is a C_1 - C_4 alkylene group optionally substituted with C_1 - C_2 alkyl;

Z is a bond, or C_1 - C_3 alkylene optionally substituted with C_1 - C_2 alkyl;

R_a is hydrogen, C_1 - C_6 alkyl, chloro, bromo, fluoro, or trifluoromethyl;

20 R_1 is hydrogen, C_1 - C_6 alkyl, halogen, preferably chloro or fluoro, or phenyl optionally substituted with up to three groups independently selected from halogen, C_1 - C_6


alkyl, C_1-C_6 alkoxy, amino, and mono- or $di(C_1-C_6)$ alkylamino;

R_2 , R_3 , R_4 and R_5 are each independently hydrogen, halogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens), nitro, OR_7 , SR_7 , $S(O)R_7$, $S(O)_2N(R_7)_2$, $C(O)N(R_7)_2$ or $N(R_7)_2$, wherein each R_7 is independently hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, amino, and mono- or $di(C_1-C_6)$ alkylamino;

phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-, 3-, or 4-pyridyl, each of which phenyl or heteroaryl is optionally substituted with up to three groups independently selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, amino, and mono- or $di(C_1-C_6)$ alkylamino;

phenoxy where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, amino, and mono- or $di(C_1-C_6)$ alkylamino; or

a group of the formula



where

J is a bond, CH₂, oxygen, or nitrogen; and

each r is independently 2 or 3;

R₆ is hydroxy, C₁-C₆ alkoxy, or -O⁻M⁺ where M⁺ is a cation forming a pharmaceutically acceptable salt; and

5 R₁₁, R₁₂, R₁₃ and R₁₄ are independently hydrogen, halogen, nitro, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, trifluoromethyl, trifluoromethoxy, C₁-C₆ alkylsulfinyl, or C₁-C₆ alkylsulfonyl.

In preferred compounds of Formula III, the R₂, R₃, R₄ and
10 R₅ substituents, in combination, represent one of bromo, cyano or nitro, one or two of fluoro, chloro, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, or trifluoromethyl, or two fluoro or two methyl with one hydroxy or one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one methyl, or three fluoro groups.
15 Particularly preferred R₂, R₃, R₄ and R₅ substituents are, independently, fluorine, chlorine, nitro, and trifluoromethyl.

In preferred compounds of Formulas II and III, A is preferably methylene, methylene substituted with a methyl group, or ethylene.

20 Preferred compounds according to Formula II above include those wherein R₈ is fluorine, R₉ is hydrogen and R₁₀ is bromine or those wherein R₈ and R₁₀ are hydrogens and R₉ is nitro.

Preferred compounds of Formula III above are those wherein the benzothiazole moiety is substituted with nitro, one, two,
25 or three of fluoro, one or two of chloro, or at least one

trifluoromethyl group. More preferred compounds of Formula II are those where A is methylene, R₁ is hydrogen or methyl, Z is a bond, and R₆ is hydroxy or C₁-C₆ alkoxy.

Still more preferred compounds of Formula II are those
5 wherein R₁₁, R₁₂ and R₁₄ are fluorines and R₁₃ is hydrogen. Other more preferred compounds of Formula II are those where R_a is methyl or hydrogen, Z is methylene or, more preferably, a bond, A is CHF or C₁ or C₂ alkylene, preferably methylene, R₁ is methyl or hydrogen, and R₁₁, R₁₂ and R₁₄ are halogens or C₁-C₃
10 alkyl. Still other more preferred compounds of Formula III are those where R_a is methyl or hydrogen, Z is methylene or, more preferably, a bond, A is CHF or C₁ or C₂ alkylene, R₁ is methyl or hydrogen, and R₁₁, R₁₂ and R₁₄ are fluorines or chlorines.

Particularly preferred compounds of Formula I are those
15 where R₃ and R₄ are independently hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or halogen, and R_a is methyl or hydrogen, Z is a bond, A is methylene, methyl substituted methylene, or ethylene, R₁ is methyl or hydrogen, and R₁₁, R₁₂ and R₁₄ are fluorines or chlorines.

20

The term "prodrug group" denotes a moiety that is converted in vivo into the active compound of formula I wherein R₆ is hydroxy. Such groups are generally known in the art and include ester forming groups, to form an ester prodrug, such as
25 benzyloxy, di(C₁-C₆)alkylaminoethyloxy, acetoxymethyl,

pivaloyloxymethyl, phthalidoyl, ethoxycarbonyloxyethyl, 5-methyl-2-oxo-1,3-dioxol-4-yl methyl, and (C₁-C₆)alkoxy optionally substituted by N-morpholino and amide-forming groups such as di(C₁-C₆)alkylamino. Preferred prodrug groups include 5 hydroxy, C₁-C₆ alkoxy, and O⁺M⁺ where M⁺ represents a cation. Preferred cations include sodium, potassium, and ammonium. Other cations include magnesium and calcium. Further preferred prodrug groups include O⁺M⁺⁺ where M⁺⁺ is a divalent cation such as magnesium or calcium.

10 In certain situations, compounds of Formula I may contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. These compounds can be, for example, racemates or optically active forms. In these situations, the single enantiomers, i.e., optically active 15 forms, can be obtained by asymmetric synthesis or by resolution of the racemates. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography, using, for example a chiral HPLC column.


20 Representative compounds of the present invention include the pharmaceutically acceptable acid addition salts of compounds where R₆ includes basic nitrogen atom, i.e., an alkylamino or morpholino group. In addition, if the compound or prodrug of the invention is obtained as an acid addition 25 salt, the free base can be obtained by basifying a solution of

the acid salt. Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds.

Non-toxic pharmaceutical salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic, alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{ACOOH}$ where n is 0-4, and the like. Non-toxic pharmaceutical base addition salts include salts of bases such as sodium, potassium, calcium, ammonium, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

As used herein, the terms 2-benzothiazolyl and benzothiazol-2-yl are synonymous.

Representative groups of the formula



include those where J is oxygen and each r is 2 (morpholinyl), J is nitrogen and each r is 2 (piperazinyl) or one r is 2 and the other 3 (homopiperazinyl), or J is CH_2 and each r is 2 (piperidinyl) or one r is 2 and the other 3 (homopiperidinyl).

Preferred groups of this formula are morpholinyl and piperazinyl.

The heterocyclic 5-membered ring having one to three nitrogen atoms, one of which may be replaced by oxygen or sulfur includes imidazolyl, oxazolyl, thiazolyl, pyrazolyl, oxadiazolyl, thiadiazolyl, and triazolyl.

The heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur includes triazinyl, pyrimidyl, pyridazinyl, oxazinyl and triazinyl.

The heterocyclic ring may be condensed with benzo so that said ring is attached at two neighboring carbon atoms to form a phenyl group. Such benzoheterocyclic ring may be attached to Z either through the heterocyclic group or through the benzo group of the benzoheterocyclic ring. Specific wherein said heterocyclic ring is condensed with a benzo include benzoxazolyl, quinazolin-2-yl, 2-benzimidazolyl, quinazolin-4-yl and benzothiazolyl. The oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms include positional isomers such as oxazolo[4,5-b]pyridine-2-yl, thiazolo[4,5-b]pyridine-2-yl, oxazolo[4,5-c]pyridine-2-yl, thiazolo[4,5-c]pyridine-2-yl, oxazolo[5,4-b]pyridine-2-yl, thiazolo[5,4-b]pyridine-2-yl, oxazolo[5,4-c]pyridine-2-yl, and thiazolo[5,4-c]pyridine-2-yl.

The following compounds of the invention are provided to give the reader an understanding of the compounds encompassed by the invention:

- 5 3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
5-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
2-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
10 5-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
7-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
6-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-
15 acetic acid
5-benzyloxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
6-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
20 5-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
6-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid
3-methyl(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-2
25 propionic acid

3-methyl (4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-3

propionic acid

3-(5-trifluoromethylbenzothiazol-2-yl)methyl-indole-N-acetic
acid

5 5-methyl-3-(5-trifluoromethylbenzothiazol-2-yl)methyl-indole-N-
acetic acid

3-(3-nitrophenyl)methyl-indole-N-acetic Acid

10 The above compounds, further described in the Examples and
other description of the invention below, are illustrative but
are not meant to limit in any way the scope of the contemplated
compounds according to the present invention.

The compounds of the invention are administered to a
15 patient or subject in need of treatment either alone or in
combination with other compounds having similar or different
biological activities. For example, the compounds of the
invention may be administered in a combination therapy, i.e.,
either simultaneously in single or separate dosage forms or in
20 separate dosage forms within hours or days of each other.
Examples of such combination therapies include administering
the compounds of Formula I with other agents used to treat
hyperglycemia, hyperlipidemia, and diabetic complications.

Suitable compounds for use in combination therapy include

25 For Hyperglycemia:

Insulin

Metformin

Troglitazone

Pioglitazone

5 Rosiglitazone

Darglitazone

Sulfonylureass such as glipizide and glimepiride

Repaglinide

alpha-glucosidase inhibitors such as acarbose, miglitol

10

For Diabetic complications:

ACE inhibitors: Captopril, lisinopril

Angiotensin II receptor antagonists (AT1-receptor) such as
candesartan, losartan, irbesartan, and valsartan

15 MMP inhibitors

Protein kinase C inhibitors

For Antihyperlipidemia:

Statins such as Atorvastatin, simvastatin, pravastatin,
20 fluvastatin, lovastatin, cerivastatin

Fibrates such as Fenofibrate, bezafibrate, ciprofibrate,
gemfibrozil

The compounds of general Formula I may be administered
25 orally, topically, parenterally, by inhalation or spray or

rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, 5 intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic 10 pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, 15 dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain 20 one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically 25 acceptable excipients which are suitable for the manufacture of

tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding
5 agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action
10 over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an
15 inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in
20 admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing
25 or wetting agents may be a naturally-occurring phosphatide, for

example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or
5 condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The
10 aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the
15 active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and
20 flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the
25 active ingredient in admixture with a dispersing or wetting

agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring
5 agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these.
10 Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation
15 products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or
20 sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those
25 suitable dispersing or wetting agents and suspending agents

which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among
5 the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono-or
10 diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared
15 by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

20 Compounds of general Formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be
25 dissolved in the vehicle.

Dosage levels on the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 1000 mg of an active ingredient.


It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The compounds of the present invention may be prepared by use of known chemical reactions and procedures. General methods for synthesizing the compounds are presented below. It is understood that the nature of the substituents required for the desired target compound often determines the preferred method of synthesis. All variable groups of these methods are as described in the generic description if they are not specifically defined below. More detailed procedures for

particular examples are presented below in the experimental section.

Methods of Preparation

- 5 The compounds of the invention where Ar is benzothiazolyl can be conveniently prepared from a substituted indole moiety using general Scheme A set forth below.



10

Scheme A

Treatment of a nitrile indole IV with a strong base such as, for example, sodium hydride, butyl lithium or sodium tert-

butoxide, in a polar aprotic solvent such as acetonitrile, tetrahydrofuran or *N,N*-dimethylformamide followed by an treatment with an alkylating agent, e.g., ethyl or *tert*-butyl bromoacetate, provides the desired *N*-alkylated product V.

5 Alternatively, phase transfer catalysis can be used in a biphasic solvent system. A general review of such alkylations can be found in Sundberg, R. J. *Indoles*; Chapter 11, Academic Press Inc., San Diego, CA, 1996. Condensation with a suitable 2-amino thiophenol hydrochloride salt VI provides

10 benzothiazole intermediate VII. These reactions are most often carried out in an alcohol solvents at elevated temperatures; however, other solvents like *N,N*-dimethylformamide and *N*-methylpyrrolidone can be used or the reactions can be carried out in the absence of solvents altogether. The scope of the


15 reaction conditions useful for this transformation have been described previously (U.S. Pat. No. 5,700,819). General methods for the preparation of various substituted 2-amino thiophenols are also well known (*J. Med. Chem.* 1991, 34, 108 and *Chem. Pharm. Bull.* 1994, 42, 1264). In general, the best

20 method of synthesis is determined by such factors as availability of starting materials and ease of synthesis. Deprotection of the alkanoic acid moiety VII can be carried out by methods common to those skilled in the art to result in compounds of Formula III. The method used in the deprotection

25 depends on the type of protecting group. A description of such

protecting groups and methods for deprotecting them may be found in: *Protective Groups in Organic Synthesis*, Second Edition, T. W. Green and P. G. M. Wuts, John Wiley and Sons, New York, 1991. When a methyl or ethyl ester is used, an aqueous sodium hydroxide solution in ethanol or dimethoxyethane is conveniently employed for its removal.

If not commercially available, nitrile IV can be prepared substantially as described below in Scheme B depicting the formation of 3-acetonitrile substituted indoles of Formula IV where Z is a bond. Thus, an indole moiety in a weak acid solution, for example, acetic acid in ethanol, is treated with aqueous formaldehyde and dimethyl amine in an alcohol solvent. The 3-(dimethylamino)methyl indole product can then be treated with sodium or potassium cyanide in *N,N*-dimethylformamide at elevated temperatures to provide the 3-acetonitrile substituted indole intermediate. Alternatively, an iminium salt like *N,N*-dimethylmethyleneammonium chloride can be used to prepare the 3-(dimethylamino)methyl indole intermediate.




Scheme B

The 3-(dimethylamino)methyl indole intermediate can also be converted to the 3-acetonitrile substituted indole intermediate via the trimethyl ammonium salt. The salt can be prepared by treating the gramine intermediate with an alkalating agent like methyl iodide. The trimethyl ammonium salt intermediate can then be converted to the nitrile by treatment with sodium or potassium cyanide in a solvent like *N,N*-dimethylformamide. In general, the conversion to the acetonitrile occurs under more mild conditions when the trimethyl ammonium salt is used.

Alternatively, other compounds, such as those where Z-Ar represents a wide variety of substituted heterocycles, may be prepared using the general method outlined in Scheme C. Here, substituted indole intermediates where X is an activating group like hydroxyl, halogen, dialkyl amino, trialkyl ammonium or

benzotriazole are coupled with Q-Z-Ar groups using methods well-established in indole chemistry. Examples of these methods where Q is Na or H and Z is sulfur, oxygen, nitrogen carbon or a bond are described in (A) Tidwell, J.H.; Peat, A.J.; Buchwald, S.L. *J. Org. Chem.* **1994**, *59*, 7164; (B) Bruneau, P.; Delvare, C.; Edwards, M.P.; McMillan, R.M. *J. Med. Chem.* **1991**, *34*, 1028; (C) Gan, T.; Cook, J.M. *Tetrahedron Lett.* **1997**, *38*, 1301; (D) Cerreto, F.; Villa, A.; Retico, A.; Scalzo, M. *Eur. J. Med. Chem.* **1992**, *27*, 701; (E) Majchrzak, M.W.; Zobel, J.N.; Obradovich, D.J.; *Synth. Commun.* **1997**, *27*, 3201; (F) DeLeon, C.Y.; Ganem, B. *J. Org. Chem.* **1996**, *61*, 8730; (G) Katritzky, A.R.; Toader, D; Xie, L. *J. Org. Chem.* **1996**, *61*, 7571.

It is understood that, depending on the specific chemistry used, a protecting group, P, may be required. In general, P represents groups such as acyloxy, alkyl, sulfonyl or A-COOR. The use of these general methods is illustrated in *Protective Groups in Organic Synthesis*, Second Edition, T. W. Green and P. G. M. Wuts, John Wiley and Sons, New York, 1991.




Scheme C

In general, the intermediate compounds wherein $R_{1,2}$ is aryl or heteroaryl can be synthesized by the chemistry illustrated in reaction Scheme D below. For example, treatment of the potassium salt of an optionally substituted bromoindole with
5 tert-butyllithium at low temperature in an ethereal solvent such as ether or tetrahydrofuran followed by the addition of an electrophile represents a general method for obtaining substituted indoles, as described by Rapoport, H. (*J. Org. Chem.* 1986, 51, 5106). For a discussion of a synthesis where R
10 is acyl, see *Biorg. Med. Chem. Lett.* 1999, 9, 333; where R is, thiomethyl, see *Heterocycles*, 1992, 34, 1169; and where R is cycloalkyl, see *J. Med. Chem.* 1999, 42, 526.

More specifically the addition of a trialkyl borate followed by an acidic work-up provides the desired indole
15 boronic acids (*Heterocycles*, 1992, 34, 1169). Indole boronic acids can be used in well established transition metal catalyzed coupling reactions like the Suzuki reaction to provide aryl and heteroaryl indoles. These reactions are most often carried out in a mixture of ethereal or alcohol solvents
20 with aqueous base in the presence of palladium catalyst, such as $\text{Pd}(\text{OAc})_2$, $\text{Pd}(\text{OAc})_2$ w/ PPh_3 or $\text{Pd}(\text{PPh}_3)_4$ as described in *Tetrahedron Lett.* 1998, 39, 4467, *J. Org. Chem.* 1999, 64, 1372 and *Heterocycles* 1992, 34, 1395.


Alternatively, an optionally substituted bromoindole can
25 be treated with an arylboronic acid and a palladium catalyst

to provide arylindoles in large quantities (Synlett 1994, 93). A general review of Suzuki cross-couplings between boronic acids and aryl halides can be found in Miyaura, N; Suzuki, A. Chem. Rev. 1995, 95, 2457.




Scheme D

For example, treatment of the advanced intermediate indole
10 X with an aryl or heteroaryl boronic acid using Pd-mediated
coupling conditions provides the desired aryl and heteroaryl
indole product XI as shown in scheme (E). In general the
utility of this method is determined by the ease of synthesis
of advanced intermediates of type X and the commercial
15 availability of aryl and heteroaryl boronic acids.




Scheme E

In addition, certain organometallic reactions eliminate the need for de novo construction of the indole nucleus. For example, the Stille reaction serves as a general method for the synthesis of regiocontrolled substitution of indole intermediates as described by Farina, V.; Krishnamurthy, V; Scott, W., *Organic Reactions*, 1998, 50, 1-652. As indicated in the scheme below, the indole may serve as the organotin species or the aryl halide. The stannylindole (XII), where P is a suitable protecting group such as [2-(trimethyl)ethoxy]methyl (SEM) or an alkyl substituent, is treated with a variety of partners (i.e., vinyl/allylic halides, vinyl triflates, aryl/heteroaryl halides and acyl halides) in the presence of a Pd(0)L_n catalyst to provide the desired indoles (XII) (Synnlett 1993, 771, *Helv. Chim. Acta* 1993, 76, 2356 and *J. Org. Chem.* 1994, 59, 4250). Conversely, a haloindole (XIV) is treated with a variety of tin reagents under Stille conditions to provide the desired substituted indoles (XV) as described in *Heterocycles* 1988, 27, 1585 and *Synth. Comm* 1992, 22, 1627).




A general procedure for the synthesis of intermediate compounds using amines of the formula $\text{NR}_x\text{R}_{x_2}$ (NR_xR_x in the scheme below) is given in scheme F below. In Scheme F, R_x and R_{x_2} are the same or different and represent hydrogen, $\text{C}_1\text{-C}_6$ alkyl, or R_x and R_{x_2} together represent a group of the formula:



where J and each r is as defined above for formula I.

As shown in Scheme F, nucleophilic substitution of X (X is halogen, preferably fluorine) in an aromatic system is a method often used to substitute aromatic rings with amine and ether functionalities. Both 4- and 5- fluoro-2-nitrotoluene are sufficiently activated to undergo substitution with amines in the presence of K_2CO_3 in a polar aprotic solvent such as, for example, DMSO as described in *J. Med. Chem.* 1993, 36, 2716. The Leimgruber -Batcho two-step method is a general process for the construction of the indole ring system from the appropriate o-nitrotoluene. This reaction involves the condensation of an o-nitrotoluene with *N,N*-dimethylformamide dimethyl acetal followed by a reductive cyclization under suitable conditions such as hydrogen over a palladium catalyst or Zn/HOAc as described in Sundberg, R.J. *Indoles*; Chapter 2, Academic Press

Inc., San Diego, CA, 1996. A representative description of the process can also be found in *Organic Synthesis*, 1984, 63, 214.




5

Scheme F

A general procedure for the synthesis of intermediate compounds wherein R is an aromatic, heteroaromatic or alkyl group is indicated in Scheme G below. As previously described, nucleophilic substitution of halogen, preferably fluorine, in an aromatic system is a method often used to substitute aromatic rings with amine and ether functionalities. Both 4- and 5-fluoro-2-nitrotoluene are sufficiently activated enough to undergo substitution with alcohols or phenols in the presence of K₂CO₃ in a polar aprotic solvent such as DMSO. A similar system using KOH and phenol is described in *J. Med. Chem.* 1994, 37, 1955. Alternatively, solid-liquid phase transfer catalysis (PTC) methods have been used to prepare intermediate ethers of this type as described in *Synth. Comm.*


1990, 20, 2855. The appropriately substituted o-nitrotoluene can then be converted to the appropriate indole by the Leimgruber-Batcho method previously described.




Scheme G

10 The preparation of intermediate alkoxy indole compounds wherein R is C₁-C₆ alkyl is outlined in Scheme H below. Commercially available nitrophenols can be alkylated under mild conditions with a base such as, for example, K₂CO₃ or Cs₂CO₃, in a polar aprotic solvent, e.g. CH₃CN, with a variety of

15 suitable alkyl halides. See *Synth. Comm.* 1995, 25, 1367. The alkoxy o-nitrotoluene can then be converted to the desired indole as described above.






Scheme H

Alternatively, some examples of the invention where Z is a
5 bond and Ar is a substituted heterocycle such as a thiazole; or
Z is amide and Ar is a substituted phenyl can be conveniently
prepared from an indole 3-acetic acid derivative as illustrated
in Scheme I. Using this method, the carboxylic acid moiety is
activated and coupled with an aryl amine. Some examples of
10 activating methods well-known to those skilled in the art
include formation of acid chloride, mixed anhydrides and
coupling reagents such as 1,3-dicyclohexylcarbodiinide (DCC).
A review of such method can be found in Bodanszky, M.
Principles of Peptide Synthesis; Springer-Verlag: New York,
15 1984. For the examples where Z is a bond and Ar is a
substituted benzothiazole or benzoxazole, the intermediate
amide or thioamide can be cyclized into the aromatic ring.
Examples of these types of heterocycle forming reactions are
described in Mylar, B. L. et al. *J. Med. Chem.* 1991, 34, 108.
20 In addition, the carboxylic acid can be converted to a chloro-
or bromomethyl ketone and condensed with nucleophiles like
thioamides or 2-aminothiophenols to produce thiazole or
benzothiazine derivatives. Examples of methods to prepare the
chloro- and bromomethyl ketones are illustrated in Rotella,

D.P.; Tetrahedron Lett. 1995, 36, 5453 and Albeck, A.; Persky, R.; Tetrahedron 1994, 50, 6333. Depending on the reaction conditions in a given synthetic sequence a protecting group may be required. It is also understood that the specific order of steps used in the synthesis depends on the particular example being prepared. P may represent H, A-COOH, A-COO-lower alkyl or a simple protecting group that can be removed at a late stage of the synthesis. When such a protecting group is used, the A-CO₂R₆ group can be introduced near the end of the synthesis after the Z-Ar group has been assembled. Method of introducing the Z-Ar group are similar to those already described.



Scheme I

5

Another strategy involves the synthesis of substituted indoles via an intramolecular cyclization of an aniline

nitrogen onto a substituted alkyne as shown in Scheme J.


Typical approaches utilize commercially available o-iodoaniline derivatives. When these intermediates are unavailable, the

regioselective ortho iodination of aromatic amines is used to

5 generate the required intermediate (*J. Org. Chem.* 1996, 61, 5804). For example, Iodophenyl intermediates are treated with trimethylsilylacetylene in the presence of a Pd catalyst and a Cu(I) source, such as cupric iodide, to produce o-

10 alkynylanilines. See *Heterocycles*, 1996, 43, 2471 and *J. Org. Chem.* 1997, 62, 6507. Further elaboration of the o-alkynylaniline to the desired indole can be done by a copper-mediated cyclization or a base-induced amine ring closure onto the alkyne functionality (*J. Med. Chem.* 1996, 39, 892). Alternative modifications have been made in the acetylenic

15 derivatives to generate more elaborate indole structures as described in *J. Am. Chem. Soc.* 1991, 113, 6689, *Tetrahedron Lett.* 1993, 24, 2823 and *Tetrahedron Lett.* 1993, 34, 6471.



Scheme J

20

Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention, as demonstrated by the following examples. In some cases,


protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general, the need for such protecting groups will be apparent to those skilled in the art of organic synthesis as well as the conditions necessary to attach and remove such groups.

The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.

The preparation of the compounds of the present invention is illustrated further by the following examples, which are not to be construed as limiting the invention in scope or spirit to the specific procedures and compounds described in them.

Example 1:

15 Preparation of 2-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid




2-Methyl-3-(4,5,7-Trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 2-methylindole was used instead of 5-chloroindole in part 1: 178-180°C; ¹H NMR (DMSO-d₆, 300 MHz) δ 7.75-7.62 (m, 1 H), 7.45 (d, J = 9.0 Hz, 1 H), 7.39 (d,

$J = 9.0$ Hz, 1 H), 7.08 (t, $J = 9$ Hz, 1 H), 6.99 (t, $J = 9.0$ Hz, 1 H), 5.00 (s, 2 H), 4.60 (s, 2 H), 2.38 (s, 3 H); LRMS calcd for $C_{19}H_{13}F_3N_2O_2S$: 390.0; found 391.0 ($M + 1$)⁺. Anal. Calcd for $C_{19}H_{13}F_3N_2O_2S$: C, 58.46; H, 3.36; N, 7.18; S, 8.21. Found: C, 58.47; H, 3.29, N, 7.12, S, 8.18.

Example 2:

Preparation of 5-chloro-3-(4,5,7-Trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid



5-chloroindole-3-acetonitrile :

A solution of aqueous formaldehyde (37%, 2.95 mL, 66.0 mmol) and dimethylamine (40%, 5.30 mL, 66.0 mmol) in 20 mL EtOH was cooled to 0°C. 5-Chloroindole (4.0 g, 26.4 mmol) was dissolved in a HOAc:EtOH mixture (1:1, 40 mL) and added dropwise to the reaction mixture. After stirring at this temperature for 2 h, the mixture was allowed to warm to room temperature and stir overnight. The mixture was added to a sat'd solution of $NaHCO_3$. 1 N NaOH was added until the pH was between 9-10. The resulting mixture was extracted with CH_2Cl_2 (3X). The organics were combined and washed with a sat'd aq. NaCl, dried over $MgSO_4$, filtered and concentrated in vacuo to

give 4.65 g (85%) of 5-chloro-3-[(dimethylamino)methyl] indole as a yellow powder. Without further purification, 5-chloro-3-[(dimethylamino)methyl] indole (4.65 g, 22.4 mmol) was dissolved in dimethylformamide (80 mL) at room temperature with stirring. To this was added KCN (2.18 g, 33.5 mmol) in H₂O (10 mL). The mixture was warmed to 140 °C and stirred for 14 h. H₂O was added and the mixture was extracted with EtOAc (2X). The organics were combined and washed with sat'd brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by SiO₂ flash chromatography (3:2, Heptane: EtOAc) to give 2.65 g (63%) of 5-chloroindole-3-acetonitrile. ¹H NMR (DMSO-d₆, 300 MHz) δ 11.30 (br s, 1 H), 7.63 (s, 1 H), 7.42-7.38 (m, 2 H), 7.05 (d, J = 6.0 Hz, 1 H), 5.70 (s, 2 H),


15 5-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid :

5-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid was prepared in a manner analogous to that set forth in Example 3 (parts 1-7), except 5-chloroindole-3-acetonitrile was used instead of 3-indolyl acetonitrile in part 20 5: mp 188-189 °C; ¹H NMR (DMSO-d₆, 300 MHz) δ 7.73-7.68 (m, 1 H), 7.63 (d, J = 1.8 Hz, 1 H), 7.51 (s, 1 H), 7.45 (d, J = 9.0 Hz, 1 H), 7.14 (dd, J₁ = 9.0, J₂ = 2.4 Hz, 1 H), 5.04 (s, 2 H), 4.65 (s, 2 H); LRMS calcd for C₁₈H₁₀F₃N₂O₂SCl: 410.0; found 411.0

(M + 1)⁺. Anal. Calcd for C₁₈H₁₀F₃N₂O₂SCl: C, 52.63; H, 2.45; N, 6.82; S, 7.81. Found: C, 52.56; H, 2.40, N, 6.71, S, 7.72.

Example 3:

5 Preparation of 3-(4,5,7-Trifluorobenzothiazol-2-yl)methyl-
 indole-N-acetic Acid



2,3,5,6-Tetrafluoroacetanilide:

A solution of 2,3,5,6-tetrafluoroaniline (200 g, 1.21
10 mol) in anhydrous pyridine (103 mL, 1.27 mol) was treated with
acetic anhydride (120 mL, 1.27 mol) and heated to 120 °C for 2
h. After cooling to room temperature, the solution was poured
into ice-cold water (500 mL). The resulting precipitate was
filtered, dissolved in ethyl acetate, dried over MgSO₄,
15 filtered and concentrated. The solid material was washed with
heptane (200 mL) and dried to give 2,3,5,6-
tetrafluoroacetanilide as a white crystalline solid (206 g,
82%): mp 136-137 °C; R_f 0.48 (50% ethyl acetate in heptane); ¹H
NMR (DMSO-d₆, 300 MHz) δ 10.10 (s, 1 H), 7.87-7.74 m, 1 H),
20 2.09 (s, 3 H). Anal. Calcd for C₈H₅F₄NO: C, 46.39; H, 2.43; N,
6.67. Found C, 46.35; H, 2.39; N, 6.68.

2,3,5,6-Tetrafluorothioacetanilide:

A flame-dried, 4-necked 5,000 mL round-bottomed flask was charged with phosphorous pentasulfide (198 g, 0.45 mol) and diluted with anhydrous benzene (3,000 mL, 0.34 M). 2,3,5,6-tetrafluoroacetanilide (185 g, 0.89 mol) was added in one portion and the bright yellow suspension was heated to a gentle reflux for 3 h. The solution was cooled to 0 °C and filtered. The insoluble material was washed with ether (2 x 250 mL) and the combined filtrate was extracted with 10% aq. NaOH (750 mL, 500 mL). After cooling the aqueous layer to 0 °C, it was carefully acidified with conc. HCl (pH 2-3). The precipitated product was collected by filtration and washed with water (500 mL). The yellow-orange material was dissolved in ethyl acetate (1,000 mL), dried over MgSO₄ and activated charcoal (3 g), filtered through a short pad of silica (50 g), and concentrated. The resulting solid was triturated with heptane (500 mL) and filtered to give 2,3,5,6-tetrafluorothioacetanilide (174.9 g, 88%): mp: 103-104°C; R_f 0.67 (50% ethyl acetate in heptane); ¹H NMR (DMSO-d₆, 300 MHz) δ 11.20 (s, 1 H), 8.00-7.88 (m, 1 H), 2.66 (s, 3 H). Anal. Calcd for C₈H₅F₄NS: C, 43.05; H, 2.26; N, 6.28. Found C, 43.10; H, 2.23; N, 6.19.

25 4,5,7-Trifluoro-2-methylbenzothiazole:

A flame-dried 5,000 mL round-bottomed flask equipped with over-head stirrer was charged with sodium hydride (15.9 g, 0.66 mol) and diluted with anhydrous toluene (3,000 mL, 0.2 M). The suspension was cooled to 0 °C, and treated with 2,3,5,6-tetrafluorothioacetanilide (134 g, 0.60 mol) in one portion. The solution was warmed to room temperature over 1 h, then heated to a gentle reflux. After 30 min, dimethylformamide (400 mL) was carefully added and the mixture was stirred for an additional 2 h. The solution was cooled to 0 °C and added to ice-water (2,000 mL). The solution was extracted with ethyl acetate (1,500 mL) and washed with sat'd. aq. NaCl (1,000 mL). The organic layer was concentrated to dryness, diluted with heptane and successively washed with water (300 mL) and sat'd. aq. NaCl (1,000 mL). The organic layer was dried over MgSO₄, filtered and concentrated to give 4,5,7-trifluoro-2-methylbenzothiazole (116.8 g, 96%) as a light brown solid: mp: 91-92 °C; R_f 0.56 (30% ethyl acetate in heptane); ¹H NMR (DMSO-d₆, 300 MHz) δ 7.76-7.67 (m, 1 H), 2.87 (s, 3 H); . Anal. Calcd for C₈H₄F₃NS: C, 47.29; H, 1.98; N, 6.82; S, 15.78. Found C, 47.56; H, 2.07; N, 6.82; S, 15.59.

2-Amino-3,4,6-trifluorothiophenol Hydrochloride:

A solution of 4,5,7-trifluoro-2-methylbenzothiazole (25.0 g, 123 mmol) in ethylene glycol (310 mL, 0.4 M) and 30% aq. NaOH (310 mL, 0.4 M) was degassed using a nitrogen stream then

heated to a gentle reflux (125 °C) for 3 h. The solution was cooled to 0 °C and acidified to pH 3-4 using conc. HCl (approx. 200 mL). The solution was extracted with ether (750 mL) and washed with water (200 mL). The organic layer was dried over
5 Na_2SO_4 , filtered and treated with 2,2-di-tert-butyl-4-methylphenol (0.135 g, 0.5 mol%). After concentrating to dryness, the crude product was dissolved in anhydrous methanol (200 mL) and treated with an HCl solution in 1,4-dioxane (3" mL, 4 N, 148 mmol). The resulting mixture was concentrated to
10 dryness, triturated with isopropylether (100 mL) and filtered to give 2-amino-3,4,6-trifluorothiophenol hydrochloride (19.3 g, 73%) as a light brown solid that was used without further purification. mp. 121-124 C; R_f 0.43 (30% ethyl acetate in heptane); Anal. Calcd for $\text{C}_6\text{H}_5\text{ClF}_3\text{NS}$: C, 33.42; H, 2.34; N, 6.50; S, 14.87. Found C, 33.45; H, 2.27; N, 6.48; S, 14.96.
15

3-cyanomethyl-indole-N-acetic acid, Ethyl Ester:

Under an atmosphere of nitrogen, a solution of 3-indolyl acetonitrile (25.0 g, 160 mmol) in dry acetonitrile (530 mL, 0.3 M) was treated with sodium hydride (95%, 4.2 g, 168 mmol) and stirred for 30 min. Ethyl bromoacetate (21.3 mL, 192 mmol) was added in a dropwise manner over 10 min and the solution was stirred at room temperature for 16 h. After concentrating under reduced pressure, the resulting residue was dissolved in
25 ethyl acetate and washed with sat'd. aq. NaCl. The organic

extracts were dried over MgSO_4 , filtered and concentrated. The crude product was recrystallized from heptane and ethyl acetate to give the target compound as a white crystalline solid (19 g, 49%): mp 98-99 °C; R_f 0.29 (30% ethyl acetate in heptane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.59 (dd, $J_1 = 7.8$ Hz, $J_2 = 0.6$ Hz, 1 H), 7.40 (dd, $J_1 = 8.1$ Hz, $J_2 = 0.6$ Hz, 1 H), 7.36 (s, 1 H), 7.18 (b t, $J = 7.2$ Hz, 1 H), 7.10 (b t, $J = 7.2$ Hz, 1 H), 5.12 (s, 2 H), 4.14 (q, $J = 7.2$ Hz, 2 H), 4.06, (s, 2 H), 1.20 (t, $J = 7.2$ Hz, 3 H); LRMS calcd for $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$: 242.3; found 243.0 (M + 1) $^+$. Anal. Calcd for $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$: C, 69.49; H, 5.82; N, 11.56. Found C, 69.39; H, 5.89; N, 11.59.

3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid, Ethyl Ester: Under a nitrogen atmosphere, a

15 solution of 3-acetonitrile-indole-N-acetic acid, ethyl ester (11.0 g, 45.4 mmol) in anhydrous ethanol (90 mL, 0.5 M) was treated with 2-amino-3,4,6-trifluorothiophenol hydrochloride (12.7 g, 59.0 mmol) and heated to a gentle reflux for 16 h. After cooling to room temperature, the solution was
20 concentrated under reduced pressure, diluted with ethyl acetate and washed with 2N HCl and sat'd. aq. NaCl. The organic layer was dried over MgSO_4 , filtered and concentrated. Purification by MPLC (10-50% ethyl acetate in heptane, 23 mL/min, 150 min) to give 3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-
25 acetic acid, ethyl ester (6.0 g, 36%) as a white crystalline

solid: mp 110-111 °C; R_f 0.41 (30% ethyl acetate in heptane);
 ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.74-7.66 (m, 1 H), 7.54 (d, J = 7.8
Hz, 1 H), 7.46 (s, 1 H), 7.40 (d, J = 8.1 Hz, 1 H), 7.15 (br t,
 J = 6.9 Hz, 1 H), 7.04 (br t, J = 7.8 Hz, 1 H), 5.14, s, 2 H),
5 4.66 (s, 2 H), 4.14 (q, J = 7.2 Hz, 3 H); LRMS calcd for
 $\text{C}_{20}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: 404.4; found 405.0 ($M + 1$) $^+$. Anal. Calcd for
 $\text{C}_{20}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$; C, 59.40; H, 3.74; N, 6.93; S, 7.93. Found C,
59.52; H, 3.72; N, 6.92; S, 8.04.


10 3-(4,5,7-trifluorobenzothiazol-2-yl) methyl-indole-N-acetic
acid:

A solution of give 3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid, ethyl ester (5.91 g, 14.6 mmol) in 1,2-dimethoxyethane (73 mL, 0.2 M) was cooled to 0 °C and
15 treated with aq. NaOH (1.25 N, 58 mL, 73.1 mmol) in a dropwise manner over 15 min. After the addition was complete, the solution was stirred for an additional 30 min, acidified to pH 3 with 2N HCl, and concentrated under reduced pressure. The residue was dissolved in ethyl acetate (200 mL) and washed with
20 sat'd. aq. NaCl (30 mL). The organic extract was dried over Na_2SO_4 , filtered and concentrated. The resulting material was stirred as a suspension in heptane, filtered and dried to give 3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid (5.38 g, 98%) as a pale yellow solid: mp 177-178 °C; R_f 0.44
25 (20% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ

7.74-7.65 (m, 1 H), 7.53 (d, $J = 7.5$ Hz, 1 H), 7.46 (s, 1 H), 7.40 (d, $J = 8.1$ Hz, 1 H), 7.15 (b t, $J = 6.9$ Hz, 1 H), 7.03 (b t, $J = 7.2$ Hz, 1 H), 5.03 (s, 2 H), 4.65 (s, 2 H); LRMS calcd for $C_{18}H_{11}F_3N_2O_2S$: 376.4; found 375.0 ($M - 1$). Anal. Calcd for $C_{18}H_{11}F_3N_2O_2S$: C, 57.44; H, 2.95; N, 7.44; S, 8.52. Found C, 57.58; H, 2.99; N, 7.38; S, 8.51.

Example 4:

Preparation of 5-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid



5-Methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 5-methylindole was used instead of 5-chloroindole in part 1: mp 131-133 °C; ¹H NMR (DMSO-d₆, 300 MHz) δ 7.73-7.62 (m, 1 H), 7.39 (s, 1 H), 7.30 (s, 1 H), 7.27 (d, $J = 9.0$ Hz, 1 H), 6.96 (dd, $J_1 = 9.0$ Hz, $J_2 = 2.4$ Hz, 1 H), 4.98 (s, 2 H), 4.60 (s, 2 H), 2.32 (s, 3 H); LRMS calcd for $C_{19}H_{13}F_3N_2O_2S$: 390.0; found 391.0 ($M + 1$). Anal. Calcd for $C_{19}H_{13}F_3N_2O_2S$: C, 58.46; H, 3.36; N, 7.18; S, 8.21. Found: C, 58.36; H, 3.30, N, 7.10, S, 8.20.

Example 5:

Preparation of 7-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

7-Methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 7-methylindole was used instead of 5-chloroindole in part 1: mp 216-218 °C; ¹H NMR (DMSO-d₆, 300 MHz) δ 7.73-7.63 (m, 1H), 7.36-7.32 (m, 2 H), 6.92-6.88 (m, 2 H), 5.17 (s, 2 H), 4.60 (s, 2 H), 2.55 (s, 3 H); LRMS calcd for C₁₉H₁₃F₃N₂O₂S: 390.0; found 391.0 (M + 1)⁺. Anal. Calcd for C₁₉H₁₃F₃N₂O₂S: C, 58.46; H, 3.36; N, 7.18; S, 8.21. Found: C, 58.37; H, 3.37; N, 7.11; S, 8.13.

Example 6:


15 Preparation of 6-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

6-Chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 6-chloroindole was used instead of 5-chloroindole in part 1: mp 194-195°C; ¹H NMR (DMSO-d₆, 300 MHz) δ 7.73-7.63 (m, 1 H), 7.50 (d, J = 8.4 Hz, 1 H), 7.46-7.42 (m, 2 H), 7.00 (dd, J₁ = 8.4 Hz, J₂ = 2.1 Hz, 1 H), 4.76 (s, 2 H), 4.62 (s, 2 H); LRMS calcd for C₁₈H₁₀F₃N₂O₂SCl: 410.0; found 411.0 (M + 1)⁺. Analysis calculated for C₁₈H₁₀F₃N₂O₂SCl: C,

52.63; H, 2.45; N, 6.82; S, 7.81. Found: C, 52.50; H, 2.44, N, 6.74, S, 7.69.

Example 7:

5 Preparation of 5-benzyloxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid



5-Benzyloxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 5-benzyloxyindole was used instead of 5-chloroindole in part 1: mp 165-168°C; ¹H NMR (DMSO-d₆, 300 MHz) δ 7.73-7.65 (m, 1 H) 7.40-7.30 (m, 3 H), 7.28-7.10 (m, 4 H), 7.10 (d, J = 2.4 Hz, 1 H), 6.87-6.80 (m, 1 H), 5.05 (s, 2 H), 4.95 (s, 2 H), 4.57 (s 2 H); LRMS calcd for C₂₅H₁₇F₃N₂O₂S: 482.0; found 483.0 (M + 1)⁺.

Example 8:


Preparation of 6-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

20 6-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 6-fluoroindole was used instead.

of 5-chloroindole in part 1: mp 200-203 C; ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.73-7.65 (m, 1 H), 7.53 (dd, $J_1 = 8.4$ Hz, $J_2 = 3.3$ Hz, 1 H), 7.44 (s, 1 H), 7.34 (dd, $J_1 = 10.5$ Hz, $J_2 = 2.4$ Hz, 1 H), 6.93-6.68 (m, 1 H), 5.11 (s, 2 H), 4.64 (s, 2 H); LRMS calcd for $\text{C}_{18}\text{H}_{10}\text{F}_4\text{N}_2\text{O}_2\text{S}$: 394.0; found 395 (M + 1).

Example 9:

Preparation of 5-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid



5-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 5-fluoroindole was used instead of 5-chloroindole in part 1: mp 193-195°C; ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.65 (m, 1 H), 7.51 (s, 1 H), 7.42 (br dd, $J_1 = 9.0$ Hz, $J_2 = 4.8$ Hz, 1 H), 7.34 (br dd, $J_1 = 9.9$ Hz, $J_2 = 2.4$ Hz, 1 H), 7.02-6.96 (m, 1 H), 5.03 (s, 2 H), 4.62 (s, 2 H); LRMS calcd for $\text{C}_{18}\text{H}_{10}\text{F}_4\text{N}_2\text{O}_2\text{S}$: 394.0; found 395 (M + 1).

Example 10:

Preparation of 6-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

6-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 2, except 6-methylindole was used instead of 5-chloroindole in part 1: mp 211-213°C, R_f 0.50 (10% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.72-7.63 (m, 1 H), 7.37 (d, J = 7.1 Hz, 1 H), 7.35 (s, 1 H), 7.18 (s, 1 H), 6.85 (d, J = 8.4 Hz, 1 H), 5.08 (s, 2 H), 4.60 (s, 2 H), 2.37 (s, 3 H).

Example 11:


Preparation of 3-(5-trifluoromethylbenzothiazol-2-yl)methyl-indole-N-acetic Acid

3-(5-trifluoromethylbenzothiazol-2-yl)methyl-indole-N-acetic Acid was prepared in a manner analogous to that set forth in Example 3 (parts 5-7), except 2-amino-4-(trifluoromethyl)-benzenethiol hydrochloride was used instead of 2-amino-3,4,6-trifluorothiophenol hydrochloride in part 6: mp 233-234 °C; ^1H NMR (DMSO- d_6 , 300 MHz) δ 8.29 (s, 1 H), 8.19 (br d, J = 8.1 Hz, 1 H), 7.68 (br d, J = 9.0 Hz, 1 H), 7.49 (br d, J = 6.9 Hz, 1 H), 7.41 (s, 1 H), 7.38 (br d, J = 8.4 Hz, 1 H), 7.12 (br t, J = 6.9 Hz, 1 H), 7.00 (br t, J = 6.9 Hz, 1 H), 5.01 (s, 2 H), 4.60 (s, 2 H).

Example 12:**Preparation of 5-Methyl-3-(5-Trifluoromethylbenzothiazol-2-yl)methyl-indole-N-acetic acid**

5-Methyl-3-(5-trifluoromethylbenzothiazol-2-yl)methyl-indole-N-acetic acid was prepared in a manner analogous to that set forth in Example 2, except 5-methylindole was used instead of 5-chloroindole in part 1 and, 2-amino-4-(trifluoromethyl)benzenethiol hydrochloride was used instead of 2-amino-3,4,6-trifluorothiophenol hydrochloride in part 2 (Example 3, part 6): mp 248-249°C; ¹H NMR (DMSO-d₆, 300 MHz) δ 8.27 (s, 1 H), 8.20 (d, J = 8.4 Hz, 1 H), 7.68 (d, J = 8.4 Hz, 1 H), 7.35 (s, 1 H), 7.27 (s, 1 H), 7.25 (d, J = 8.1 Hz, 1 H), 6.95 (d, J = 8.1 Hz, 1 H), 4.96 (s, 2 H), 4.57 (s, 2 H), 2.31, (s, 3 H); LRMS calcd for C₂₂H₁₅F₃N₂O₂S; found 405 (M + H).

15

Example 13:**Preparation of 3-(3-nitrophenyl)methyl-indole-N-acetic acid****Preparation of indole-N-acetic acid, ethyl ester**

Under an atmosphere of nitrogen, a solution of indole (15.0 g, 128 mmol) in dry acetonitrile (300 mL, 0.4 M) was

treated with sodium hydride (95%, 3.69 g, 153 mmol) and stirred for 30 min. Ethyl bromoacetate (17.0 mL, 153 mmol) was added in a dropwise manner over 10 min and the solution was stirred at room temperature for 16 h. After concentrating under reduced pressure, the resulting residue was dissolved in ethyl acetate and washed with sat'd. aq. NaCl. The organic extracts were dried over MgSO_4 , filtered and concentrated. The crude product was purified by flash column chromatography (50% ethyl acetate in heptane): R_f 0.25 (40% ethyl acetate in heptane) ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 7.53 (d, $J = 6.3$ Hz, 1 H), 7.38-7.31 (m, 2 H), 7.11 (br t, $J = 7.2$ Hz, 1 H), 7.02 (br t, $J = 7.2$ Hz, 1 H), 6.45-6.43 (m, 1 H), 5.10 (s, 2 H), 4.12 (q, $J = 7.2$ Hz, 2 H), 1.19 (t, $J = 7.2$ Hz, 3 H).

Preparation of 3-(3-nitrophenyl)methyl-indole-N-acetic acid, ethyl ester


Indole-N-acetic acid, ethyl ester (0.500 g, 2.50 mmol) was dissolved in 1,4-dioxane (5 mL) at room temperature with stirring. To this solution was added $\text{Ag}_2\text{CO}_3/\text{Celite}$ (50% by weight, 0.500 g, 0.9 mmol). The mixture was warmed to 90°C and maintained overnight. H_2O was added to the reaction mixture followed by extracted with EtOAc (2X). The organics were combined and washed with a sat'd brine solution, dried over MgSO_4 , filtered and concentrated in vacuo. The residue was

purified by SiO₂ flash chromatography (3:2 Heptane: EtOAc) to give 180 mg (22%) as a pale yellow oil. ¹H NMR (DMSO-d₆, 300 MHz) δ 8.10 (s, 1H), 8.02 (d, *J* = 8.1 Hz, 1 H), 7.75 (d, *J* = 7.2 Hz, 1 H), 7.59-7.57 (m, 1 H), 7.46-7.39 (m, 1 H), 7.33 (d, *J* = 8.1 Hz, 1 H), 7.20 (s, 1 H), 7.13-6.89 (m, 2 H), 5.06 (s, 2 H), 4.19 (s, 2 H), 4.13 (q, *J* = 7.2 Hz, 2 H), 1.18 (t, *J* = 7.2 Hz, 3 H).

Preparation of 3-(3-nitrophenyl)methyl-indole-N-acetic Acid

3-(3-Nitrophenyl)methyl-indole-N-acetic Acid, ethyl ester (0.175 g, 0.5 mmol) was dissolved in THF: EtOH (1:4, 5 mL) at room temperature with stirring. The mixture was cooled to 0°C and treated with 1N NaOH (1.55 mL, 1.6 mmol). The mixture was allowed to stir at this temperature for 2 h. 1 N HCl was added and the mixture extracted with EtOAc (2X). The organics were combined and washed with a sat'd brine solution, dried over MgSO₄, filtered and concentrated in vacuo. The residue was triturated with heptane and vacuum- filtered with several heptane washings to give 110 mg (69%) the desired compound as an off-white powder. mp 163-165 °C; ¹H NMR (DMSO-d₆, 300 MHz) δ 8.11 (s, 1 H), 8.03 (d, *J* = 8.1 Hz, 1 H), 7.75 (d, *J* = 8.1 Hz, 1 H), 7.53 (t, *J* = 8.1 Hz, 1 H), 7.45 (d, *J* = 8.1 Hz, 1 H), 7.33 (d, *J* = 8.4 Hz, 1 H), 7.20 (s, 1 H), 7.11 (t, *J* = 7.2 Hz, 1 H), 6.97 (t, *J* = 7.2 Hz, 1 H), 4.96 (s, 2 H), 4.18 (s, 2 H); LRMS calcd for C₁₇H₁₄N₂O₄S: 310.0; found 311 (M + 1)⁺.

Example 14

Preparation of 2-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

5

2-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid was prepared in a manner analogous to that set forth in Example 2, except that 2-phenylindole was used instead of 5-chloroindole in part 1: mp 238-239°C; R_f 0.60 (10% methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.60-7.70 (m, 1H), 7.39-7.58 (m, 7H), 7.20 (t, J = 9 Hz, 1H), 7.07 (t, J = 9 Hz, 1H), 4.80 (s, 2H), 4.45 (s, 2H); LRMS calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: 452.0; found 453.0 ($M + 1$). Anal. Calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: C, 63.71; H, 3.34; N, 6.19; S, 7.09. Found: C, 63.46; H, 3.32; N, 6.11; S, 6.96.

15

Example 15

Preparation of 5-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

20

3-cyanomethyl-5-phenyl-indole-N-acetic acid, ethyl ester

5-Bromo-3-cyanomethyl-indole-N-acetic acid, ethyl ester (1.0 g, 3.1 mmol) and phenylboronic acid (0.418 g, 3.4 mmol) were dissolved in anhydrous DME at room temperature under a

nitrogen atmosphere and treated with $\text{Pd}(\text{OAc})_2$ (2.1 mg, 0.0093 mmol) and PPh_3 (7.4 mg, 0.028 mmol). This mixture was heated to reflux and 2 M Na_2CO_3 (3.11 mL, 6.2 mmol) was added via syringe. After 12h, the mixture was cooled to room temperature and added to H_2O (50mL). The resultant mixture was extracted with EtOAc (2X, 100mL) and the organics were combined and washed with a sat'd aqueous NaCl solution, dried over MgSO_4 , filtered and concentrated in vacuo. The residue was purified by SiO_2 flash chromatography (heptane to 1:1 heptane/ EtOAc) to give the desired material as a white solid (445 mg, 45%); ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 7.64-7.74 (m, 4H), 7.39-7.44 (m, 4H), 7.29-7.34 (m, 1H), 5.20 (s, 2H), 4.15 (q, $J = 7.2$ Hz, 2H), 4.08 (s, 2H), 1.20 (t, $J = 7.2$ Hz, 3H).

15 5-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl indole-N-acetic acid

5-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid was prepared in a manner analogous to that set forth in Example 2, except that 5-phenylindole was used instead of 5-chloroindole in part 1: mp 156-159 °C; R_f 0.55 (10% methanol in chloroform); ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 7.66-7.69 (m, 4H), 7.57-7.60 (m, 1H), 7.39-7.47 (m, 3H), 7.29-7.35 (m, 2H), 5.06 (s, 2H), 4.66 (s, 2H); LRMS calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: 452.0; found 453.0 ($M + 1$) $^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: C, 63.71; H, 3.34; N, 6.19; S, 7.09. Found: C, 63.54; H, 3.32; N, 6.13; S, 7.01.

Example 16

Preparation of 6-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

5

Part 1: 6-Phenylindole

A solution of 6-bromoindole (2.0 g, 10.20 mmol) in anhydrous toluene (20mL) under a nitrogen atmosphere was treated with $\text{Pd}[\text{P}(\text{Ph}_3)]_4$ (10% mol). After stirring the mixture for 30 min., phenylboronic acid (1.87 g, 15.30 mmol) in anhydrous EtOH (10 mL) was added followed by the addition of sat'd NaHCO_3 (6mL). The bi-phasic mixture was heated to reflux for 24 h. After cooling to room temperature, the mixture was added to a sat'd brine solution and extracted with EtOAc (2X). The organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The residue was purified by flash column chromatography (1:1 CH_2Cl_2 / heptane) to give the desired material as white powder (900 mg, 45%): ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ 11.15 (br s, 1H), 7.58-7.66 (m, 4H), 7.41-7.47 (m, 2H), 7.36 (m, 1H), 7.26-7.31 (m, 2H), 6.42 (m, 1H).

20

Preparation of 6-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl indole-N-acetic acid


25

6-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid was prepared in a manner analogous to that set forth in Example 2, except that 6-phenylindole was used instead of 5-chloroindole in part 1: mp 156-159°C; R_f 0.50 (10%

methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.65-7.75 (m, 4H), 7.57-7.62 (m, 1H), 7.41-7.50 (m, 3H), 7.26-7.38 (m, 2H), 5.12 (s, 2H), 4.68 (s, 2H); LRMS calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: 452.0; found 453.0 ($M + 1$) $^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}$: C, 63.71; H, 3.34; N, 6.19; S, 7.09. Found: C, 63.46; H, 3.33; N, 6.10; S, 6.96.

Example 17

10 Preparation of 5-morpholino-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid



5-Morpholino-2-nitrotoluene

A mixture of 5-fluoro-2-nitrotoluene (5.11 g, 32.9 mmol), morpholine (4.31 mL, 49.4 mmol) and K_2CO_3 (6.83 g, 49.4 mmol) was diluted in anhydrous DMSO (80 mL) at room temperature with stirring. The mixture was heated to 80°C for 24 h. After cooling to room temperature, H_2O was added and the resultant mixture was extracted with EtOAc (3X, 50 mL). The organic layer was washed with sat'd aqueous NaCl (100 mL), dried over MgSO_4 , filtered and concentrated in vacuo. The remaining solid was triturated in heptane (200 mL) and filtered to give the desired material (7.10 g, 97%) as a yellow powder: R_f 0.40 (75%

heptane/ 25% ethyl acetate). ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.96 (d, J = 9.9 Hz, 1H), 8.85-8.88 (m, 2H), 3.70 (t, J = 5.0 Hz, 4H), 3.35 (t, J = 5.0 Hz, 4H), 2.53 (s, 3H).

5 Preparation of 5-Morpholinoindole

Under an atmosphere of nitrogen, a solution of 5-morpholinyl-2-nitrotoluene (7.0 g, 31.5 mmol) in DMF (100mL) was treated with dimethylformamide dimethyl acetal (4.81 mL, 36.2 mmol) and pyrrolidine (2.62 mL, 31.5 mL). The mixture was heated to 100°C and maintained for 12 h. After cooling, the mixture was concentrated in vacuo to give the desired intermediate as a brick-red solid.

The intermediate enamine was dissolved in EtOAc (200 mL) and added to a pre-charged Parr bottle with 10% Pd/C (600 mg) in EtOAc (40 mL). The mixture was hydrogenated on a Parr-shaker at 55 psi for 2.5 h. The catalyst was filtered through a Celite plug with several washings with EtOAc and the remaining filtrate concentrated in vacuo. The residue was purified by SiO_2 flash chromatography (1:1 Hept/EtOAc) to give 2.0 g (31% over 2 parts) of the desired indole as a cream powder: R_f 0.30 (10% methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ 10.77 (br s, 1H), 7.24 (s, 1H), 7.18-7.20 (m, 1H), 6.97 (d, J = 1.8 Hz, 1H), 6.81 (dd, J_1 = 8.7 Hz, J_2 = 2.1 Hz, 1H), 6.25 (dd, J_1 = 3.0 Hz, J_2 = 1.8 Hz, 1H), 3.7 (t, J = 4.50 Hz, 4H), 2.96 (t, J = 4.50 Hz, 4H).

Preparation of 5-morpholino-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl indole-N-acetic acid

5 5-morpholino-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl indole-N- acetic acid was prepared in a manner analogous to that set forth in Example 2, except that 5-morpholinoindole was used instead of 5-chloroindole. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.64-7.72 (m, 1H), 7.34 (s, 1H), 7.26 (d, J = 9.0 Hz, 1H), 7.06
10 (d, J = 2.4 Hz, 1H), 6.91 (dd, J₁ = 9.0 Hz, J₂ = 2.4 Hz, 1H), 4.95 (s, 2H), 4.60 (s, 2H), 3.70-3.73 (m, 4H), 2.97-3.00 (m, 4H); LRMS calcd for C₂₂H₁₈F₃N₃O₃S: 461.0; found 462 (M + 1)⁺. Anal. Calcd for C₂₂H₁₈F₃N₃O₃S·1H₂O: C, 55.11; H, 4.20; N, 8.76; S, 6.69. Found: C, 55.11; H, 4.05; N, 8.57; S, 6.50.

Example 18

Preparation of 6-morpholino-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl-indole-N-acetic acid

20 Preparation of 4-Morpholino-2-nitrotoluene

A mixture of 4-fluoro-2-nitrotoluene (15.34 g, 98.9 mmol), morpholine (12.94 mL, 49.4 mmol) and K₂CO₃ (6.83 g, 148.3 mmol) were diluted in anhydrous DMSO (250 mL) at room temperature with stirring. The mixture was heated to 120°C for 24 h.
25 After cooling to room temperature, H₂O was added and the resultant mixture was extracted with EtOAc (3X, 75 mL). The organic layer was washed with sat'd brine (100 mL), dried over

MgSO₄, filtered and concentrated in vacuo. The remaining solid was triturated in heptane (200 mL) and filtered to give the desired material (8.00 g, 36.4%) as a yellow powder: R_f 0.40 (25% ethyl acetate in heptane). ¹H NMR (DMSO-d₆, 300 MHz) δ 7.40 (d, J = 2.7 Hz, 1H), 7.30 (d, J = 8.7 Hz, 1H), 7.20 (dd, J₁ = 8.7 Hz, J₂ = 2.7 Hz, 1H), 3.70 (t, J = 4.8 Hz, 4H), 3.35 (t, J = 4.8 Hz, 4H), 2.36 (s, 3H).

Preparation of 6-Morpholinoindole

Under an atmosphere of nitrogen, a solution of 4-morpholino-2-nitrotoluene (7.1 g, 31.9 mmol) in DMF (100 mL) was treated with dimethylformamide dimethyl acetal (4.92 mL, 37.1 mmol) and pyrrolidine (2.67 mL, 31.9 mL). The mixture was heated to 100°C and maintained for 12 h. After cooling, the mixture was concentrated in vacuo to give the desired intermediate as a brick-red solid. The crude intermediate was dissolved in glacial HOAc (250 mL) and warmed to 85°C. Zn (18.17 g, 0.278 mol) was added to the solution portionwise over 30 min. The mixture was heated for 4h. After cooling to room temperature, the mixture was neutralized with sat'd NaHCO₃ and extracted with Et₂O (3X, 300 mL). The combined organics were washed with sat'd brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by SiO₂ flash chromatography (heptane to 2:1 heptane/EtOAc) to give the desired material as a white crystalline powder (1.0 g, 11% over


2 parts): R_f 0.50 (2:1 Heptane/EtOAc); ¹H NMR (DMSO)-d₆, 300
MHz) δ 10.73 (br s, 1H), 7.35 (d, J = 8.4 Hz, 1H), 7.11 (d, J =
2.4 Hz, 1H), 6.80 (s, 1H), 6.73 (dd, J₁ = 8.4 Hz, J₂ = 2.4 Hz,
1H), 6.25 (d, J = 2.4 Hz, 1H), 3.72 (t, J = 4.8 Hz, 4H), 3.02
5 (t, J = 4.8 Hz, 1H).

Preparation of 6-morpholino-3-(4,5,7-trifluorobenzothiazol-2-
yl)methyl indole-N-acetic acid

10 6-morpholino-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl
indole-N-acetic acid was prepared in a manner analogous to that
set forth in Example 2, except that 6-morpholinoindole was used
instead of 5-chloroindole in part 1: mp 178-180°C; ¹H NMR
(DMSO-d₆, 300 MHz) δ 7.66-7.72 (m, 1H), 7.37 (d, J = 8.4 Hz,
15 1H), 7.29 (s, 1H), 7.06 (d, J = 2.4 Hz, 1H), 6.84 (d, J = 8.4
Hz, 1H), 4.96 (s, 2H), 4.58 (s, 2H), 3.37-3.75 (m, 4H), 3.09-
3.13 (m, 4H); LRMS calcd for C₂₂H₁₈F₃N₃O₃S: 461.0; found 462
(M+1)⁺. Anal. Calcd for C₂₂H₁₈F₃N₃O₃S CH₂Cl₂ 0.50H₂O: C, 49.74; H,
3.72; N, 7.57; S, 5.77 Found C, 49.73; H, 3.36; N, 7.69; S,
20 5.58

Example 19

Preparation of 5-phenoxy-3-(4,5,7-trifluorobenzothiazol-2-
yl)methyl-indole-N-acetic acid



5-Phenoxy-2-nitrotoluene

A solution of phenol (12.16 g, 0.129 mol) in anhydrous
5 DMSO was treated with K_2CO_3 (17.88 g, 0.129 mol) and stirred at
room temperature for 15 min. 5-Fluoro-2-nitrotoluene (13.38 g,
0.086 mol) was added to the solution via syringe. The resultant
mixture was heated to 80°C for 12 h. After cooling to room
temperature, the mixture was poured into H_2O (100mL). After
10 extraction with EtOAc (2X, 100mL), the organics were combined
and washed with a sat'd brine solution, dried over $MgSO_4$,
filtered and concentrated in vacuo. The residue was purified by
flash column chromatography (heptane to 8:1 heptane/ EtOAc) to
give the desired material as a yellow crystalline solid (12.50
15 g, 63%): R_f 0.60 (85% heptane/ 15% EtOAc); 1H NMR ($DMSO-d_6$, 300
MHz) δ 8.05 (d, J = 9.0 Hz, 1H), 7.44-7.47 (m, 2H), 7.23-7.29
(m, 1H), 7.12-7.16 (m, 2H), 7.04 (d, J = 2.7 Hz, 1H), 6.90 (dd,
 J_1 = 9.0 Hz, J_2 = 2.7 Hz, 1H), 2.51 (s, 3H).

20 5-Phenoxyindole

A solution of 5-phenoxy-2-nitrotoluene (10.03 g, 0.0428
mol) in anhydrous DMF was treated with *N,N*-dimethylformamide

dimethyl diacetal (6.73 mL, 0.0508 mol) and pyrrolidine (3.63 mL, 0.0438 mol) and heated to 110 C for 2.5 h. After cooling to room temperature, the mixture was diluted with EtOAc (500 mL) and washed H₂O (500 mL). The organics were dried over MgSO₄,
5 filtered and concentrated in vacuo. The crude intermediate was dissolved in glacial HOAc (250 mL) and warmed to 85°C. Zn (24.62 g, 0.377 mol) was added to the solution portion wise over 30 min. The mixture was heated for 4h. After cooling to room temperature, the mixture was neutralized with sat'd NaHCO₃,
10 and extracted with Et₂O (3X, 300 mL). The combined organics were washed with sat'd brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by SiO₂ flash chromatography (heptane to 2:1 heptane/ EtOAc) to give the desired material as a white crystalline powder (3.1 g, 34% over
15 2 parts): R_f 0.50 (2:1 Heptane/ EtOAc); ¹H NMR (DMSO-d₆, 300 MHz) δ 11.12 (br s, 1H), 7.48 (s, 1H), 7.30-7.38 (m, 1H), 7.25-7.29 (m, 2H), 7.17 (d, J = 2.7 Hz, 1H), 6.89-7.02 (m, 1H), 6.86-6.88 (m, 2H), 6.80 (dd, J₁ = 8.7 Hz, J₂ = 2.4 Hz, 1H), 6.37 (m, 1H).

20

Preparation of 5-phenoxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl indole-N-acetic acid

5-phenoxy-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl
indole-N- acetic acid was prepared in a manner analogous to
25 that set forth in Example 2, except that 5-phenoxyindole was

used instead of 5-chloroindole in part 1: mp 128-130°C; R_f 0.45 (10% methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.65-7.70 (m, 1H), 7.47 (s, 1H), 7.42 (d, J = 8.4 Hz, 1H), 7.21-7.27 (m, 3H), 6.98 (m, 1H), 6.83-6.90 (m, 3H), 5.02 (s, 2H), 4.60 (s, 2H); LRMS calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_3\text{S}$: 468.0; found 467.0 ($M - 1$) $^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_3\text{S}$: C, 55.11; H, 4.20; N, 8.76; S, 6.69. Found: C, 55.11; H, 4.05; N, 8.57; S, 6.50.

Example 20

Preparation of 7-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid

7-Fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl indole-N- acetic acid was prepared in a manner analogous to that set forth in Example 2, except that 7-fluoroindole was used instead of 5-chloroindole in part 1: mp 194-196°C; R_f 0.60 (10% methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.67-7.73 (m, 1H), 7.46 (s, 1H), 7.35 (d, J = 7.2 Hz, 1H), 6.89-6.99 (m, 2H), 5.06 (s, 2H), 4.64 (s, 2H); LRMS calcd for $\text{C}_{18}\text{H}_{10}\text{F}_4\text{N}_2\text{O}_2\text{S} \cdot \text{H}_2\text{O}$: C, 50.23; H, 3.28; N, 6.51; S, 7.45. Found C, 50.70; H, 2.52; N, 6.60; S, 7.57. 394.0; found 395.0 ($M + 1$) $^+$. Anal. Calcd for $\text{C}_{18}\text{H}_{10}\text{F}_4\text{N}_2\text{O}_2\text{S}$

Example 21

Preparation of 7-bromo-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid


7-bromo-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl
indole-N-acetic acid was prepared in a manner analogous to that
set forth in Example 2, except that 7-bromoindole was used
instead of 5-chloroindole in part 1: mp 228-230°C; R_f 0.40 (10%
5 methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.65-7.74
(m, 1H), 7.57 (d, $J = 7.8$ Hz, 1H), 7.49 (s, 1H), 7.32 (d, $J =$
7.8 Hz, 1H), 6.94 (t, $J = 7.8$ Hz, 1H), 5.29 (s, 2H), 4.65 (s,
2H); LRMS calcd for $\text{C}_{18}\text{H}_{10}\text{F}_3\text{N}_2\text{O}_2\text{SBr}$: 454.0 for (^{79}Br and 456.0 for
 ^{81}Br); found 453.0 ($M - 1$) $^-$ and 455.0 ($M - 1$) $^-$. Anal Calcd for
10 $\text{C}_{18}\text{H}_{10}\text{F}_3\text{N}_2\text{O}_2\text{SBr}$: C, 47.49; H, 2.21; N, 6.15; S, 7.04. Found: C,
47.65; H, 2.27; N, 6.15; S, 6.98.

Example 22

Preparation of 7-chloro-3-(4,5,7-trifluorobenzothiazol-2-
15 yl)methyl-indole-N-acetic acid

7-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl
indole-N-acetic acid was prepared in a manner analogous to
that set forth in Example 2, except that 7-chloroindole was
used instead of 5-chloroindole in part 1: mp 228-230°C; R_f 0.38
20 (10% methanol in chloroform); ^1H NMR (DMSO- d_6 , 300 MHz) δ
7.62-7.73 (m, 1H), 7.52 (d, $J = 7.5$ Hz, 1H), 7.49 (s, 1H), 7.15
(d, $J = 7.5$ Hz, 1H), 7.00 (t, $J = 7.5$ Hz, 1H), 5.25 (s, 2H),
4.65 (s, 2H); LRMS calcd for $\text{C}_{18}\text{H}_{10}\text{F}_3\text{N}_2\text{O}_2\text{SCl}$: 410.0; found 409.0
($M - 1$) $^-$. Anal. Calcd for $\text{C}_{18}\text{H}_{10}\text{F}_3\text{N}_2\text{O}_2\text{SCl}$: C, 52.63; H, 2.45; N,
25 6.82; S, 7.81. Found: C, 52.60; H, 2.54; N, 6.66; S, 7.59.

Example 23

3-[5-Fluorbenzothiazole-2-yl]methyl-indole-N-acetic Acid

5 3-[5-fluorbenzothiazole-2-yl]methyl-indole-N-acetic acid
was prepared in a manner analogous to that set forth in Example
3, except 2-amino-4-fluorothiophenol hydrochloride was used
instead of 2-amino-4,5,7-trifluorothiophenol hydrochloride in
part 6: mp 208°C (decomp); R_f 0.10 (10% methanol in
10 dichloromethane) ^1H NMR (DMSO- d_6 , 300 MHz) δ 12.91 (s, 1 H), 7.98
(dd, J = 8.9, 5.6 Hz: 1 H), 7.78 (dd, J = 10.0, 2.6 Hz, 1 H),
7.50 (d, J = 7.8 Hz, 1 H), 7.40 (s, 1 H), 7.37 (d, J = 7.8 Hz,
1 H), 7.26 (dt, J = 8.9, 2.4 Hz, 1 H), 7.13 (t, J = 7.8 Hz, 1
H), 7.01 (t, J = 7.8 Hz, 1 H), 5.01 (s, 2 H), 4.56 (s, 2 H);
15 LRMS m/z 341.0 ($M + 1$) $^+$, 339.0 ($M - 1$) $^-$. Anal. Calcd for
 $\text{C}_{18}\text{H}_{13}\text{FN}_2\text{O}_2\text{S}$: C, 63.52; H, 3.85; N, 8.23; S, 9.42; Found: C,
63.40; H, 3.80; N, 8.37; S, 9.43.

Example 24

20 3-[6-Fluorbenzothiazole-2-yl]methyl-indole-N-acetic Acid

3-[6-fluorbenzothiazole-2-yl]methyl-indole-N-acetic acid
was prepared in a manner analogous to that set forth in Example


3, except 2-amino-5-fluorothiophenol hydrochloride was used instead of 2-amino-4,5,7-trifluorothiophenol hydrochloride in part 6: mp 203°C (decomp) R_f 0.13 (10% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 12.91 (s, 1 H), 7.95 (dd, J = 8.9, 5.0 Hz, 1 H), 7.86 (dd, J = 8.8, 2.8 Hz, 1 H), 7.50 (d, J = 7.5 Hz, 1 H), 7.40-7.35 (m, 2 H), 7.32 (dt, J = 8.9, 2.7 Hz, 1 H), 7.13 (t, J = 7.6 Hz, 1 H), 7.00 (t, J = 7.6 Hz, 1 H), 5.01 (s, 2 H), 4.54 (s, 2 H); LRMS m/z 341.0 ($M + 1$) $^+$, 339.0 ($M - 1$). Anal. Calcd for $\text{C}_{18}\text{H}_{13}\text{FN}_2\text{O}_2\text{S}$: C, 63.52; H, 3.85; N, 8.23; S, 9.42. Found: C, 63.52; H, 3.86; N, 8.35; S, 9.53.

The compounds of Examples 25-32 were prepared essentially according to the procedures set forth above in examples 1 and/or 2 with appropriate substitution of starting materials.

15

Example 25

3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-2-propionic acid




mp 176-177°C; R_f 0.34 (20% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.60-7.73 (m, 1H), 7.60 (s, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.44 (d, J = 8.1 Hz, 1H), t, J = 7.5 Hz, 1H),

7.02 (t, $J=7.5$ Hz, 1H), 5.35 (q, $J= 8.1$ Hz, 1H), 4.64 (s, 2H),
1.72 (d, $J = 8.1$ Hz, 3H); LRMS calcd for $C_{19}H_{13}F_3N_2O_2S$: 390.0;
Found 391.0 ($M + 1$)⁺. Anal. Calcd for $C_{19}H_{13}F_3N_2O_2SH_2O$: C, 55.88;
H, 3.70; N, 6.86; S, 7.85 Found: C, 56.09; H, 3.31; N, 6.89; S,
5 7.99.

Example 26

3-(4-5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-3-propionic
acid



10

mp 200-201°C; R_f 0.50 (20% methanol in dichloromethane); 1H NMR
(DMSO- d_6 , 300 MHz) δ 7.63-7.71 (m, 1H), 7.51 (s, 1H), 7.47 (d,
 $J = 3.0$ Hz, 2H), 7.14 (t, $J = 7.5$ Hz, 1H), 7.00 (t, $J= 7.5$ Hz,
1H), 4.61 (s, 2H), 4.39 (t, $J = 6.6$ Hz, 2H), 2.75 (t, $J = 6.6$
15 Hz, 2H); LRMS calcd for $C_{19}H_{13}F_3N_2O_2S$: 390.0; Found 391.0 ($M + 1$)⁺.
Anal Calcd for $C_{19}H_{13}F_3N_2O_2S$: C, 58.46; H, 3.36; N, 7.18; S, 8.21
Found: C, 58.63; H, 3.40; N, 7.20; S, 8.30.

Example 27

20 Preparation of 6-Bromo-3-(5-trifluoromethylbenzothiazol-2-
yl)methyl-indole-N-acetic acid: mp 265-267°C; R_f 0.19 (20%


methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 8.28 (s, 1H), 8.22 (d, J = 8.7 Hz, 1H), 7.67-7.69 (m, 2H), 7.43-7.47 (m, 2H), 7.14 (d, J = 9.0 Hz, 1H), 5.04 (s, 2H), 4.61 (s, 2H); LRMS calcd for $\text{C}_{19}\text{H}_{12}\text{F}_3\text{N}_2\text{O}_2\text{SBr}$: 469.0; Found 469.0 ($M + 1$) $^+$ for Br = 79. Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{F}_3\text{N}_2\text{O}_2\text{SBr}$: C, 48.63; H, 2.58; N, 5.97; S, 6.83. Found: C, 48.60; H, 2.63; N, 5.88; S, 6.91.

Example 28

6-Methoxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-
10 acetic acid: mp 118-120°C; R_f 0.27 (20% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.63-7.73 (m, 1H), 7.39 (s, 1H), 7.28 (d, J = 8.7 Hz, 1H), 7.07 (s, 1H), 6.78 (d, J = 8.7 Hz, 1H), 4.97 (s, 2H), 4.61 (s, 2H); 3.07 (s, 3H); LRMS calcd for $\text{C}_{19}\text{H}_{13}\text{F}_3\text{N}_2\text{O}_3\text{S}$: 406.0; Found 407.0 ($M + 1$) $^+$. Anal. Calcd for $\text{C}_{19}\text{H}_{13}\text{F}_3\text{N}_2\text{O}_3\text{SH}_2\text{O}$: C, 53.77; H, 3.56; N, 6.60; S, 7.56 Found:
15 C, 53.87; H, 3.56; N, 6.67; S, 7.67.

Example 29


4-Chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-
20 acetic acid



mp 203-206 °C; R_f 0.24 (20% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.63-7.71 (m, 1H), 7.57 (s, 1H), 7.33 (d, J = 9.0 Hz, 1H), 7.12 (dd, J ($_1$) = 9.0, J ($_2$) = 7.8 Hz, 1H), 7.03 (d, J = 7.8 Hz, 1H), 5.08 (s, 2H), 4.78 (s, 2H); LRMS calcd for $\text{C}_{18}\text{H}_{10}\text{F}_3\text{N}_2\text{O}_2\text{SCl}$: 410.0; Found 411.0 ($\text{M}+1$) $^+$ and 409.0 ($\text{M}-1$) $^-$.

Example 30

5-Methoxy-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl-indole-N-
 10 acetic acid



mp 165-167 °C; R_f 0.37 (20% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.61-7.70 (m, 1H), 7.35 (d, J = 9.0 Hz, 1H), 7.26 (s, 1H), 6.90 (s, 1H), 6.64 (d, J = 9.0 Hz, 1H),
 15 4.79 (s, 2H); 4.56 (s, 2H), 3.72 (s, 3H); LRMS calcd for $\text{C}_{16}\text{H}_{13}\text{F}_3\text{N}_2\text{O}_2\text{S}$: 406.0; Found 407.0 ($\text{M}+1$) $^+$ and 405.0 ($\text{M}-1$) $^-$.

Example 31


5-Bromo-3-(4,5,7-trifluorobenzothiazol-2-yl) methyl-indole-N-
 20 acetic acid: mp 209-294 °C; R_f 0.18 (20% methanol in dichloromethane); ^1H NMR (DMSO- d_6 , 300 MHz) δ 7.78 (d, J = 1.8

Hz, 1H), 7.65-7.73 (m, 1H), 7.49 (s, 1H), 7.61 (d, J = 9.0 Hz, 1H), 7.25 (dd, J (1) = 9.0 Hz, J (2) = 1.8 Hz, 1H), 5.04 (s, 2H); 4.64 (s, 2H); LRMS calcd for C₁₈H₁₀F₃N₂O₂SBr: 455.0; Found 455.0 (M+1)⁺ for Br 79 and 457 (M+1)⁺ for Br 81.

5

Example 32

3-(6-chlorobenzothiazol-2-yl) methyl-indole-N-acetic acid



10

Example 33

Representative compounds of the invention were tested for their potency, selectivity and efficacy as inhibitors of human aldose reductase. The potency or aldose reductase inhibiting effects of the compounds were tested using methods similar to those described by Butera et al. in *J. Med. Chem.* 1989, 32, 757. Using this assay, the concentrations required to inhibit human aldose reductase (hALR2) activity by 50% (IC₅₀) were determined.

In a second assay, a number of the same compounds were tested for their ability to inhibit aldehyde reductase (hALR1), a structurally related enzyme. The test methods employed were essentially those described by Ishii, et al., *J. Med. Chem.*

20

1996 39: 1924. Using this assay, the concentrations required to inhibit human aldehyde reductase activity by 50% (IC₅₀) were determined.

From these data, the hALR1 / hALR2 ratios were determined.

5 Since high potency of test compounds as inhibitors of aldose reductase is desirable, low hALR2 IC₅₀ values are sought. On the other hand, high potency of test compounds as inhibitors of aldehyde reductase is undesirable, and high hALR1 IC₅₀s values are sought. Accordingly, the hALR1 / hALR2 ratio is used to
10 determine the selectivity of the test compounds. The importance of this selectivity is described in Kotani, et al., *J. Med. Chem.* 40: 684, 1997.

The results of all these tests are combined and illustrated in Table 1.

15

Example #	hALR2 (IC ₅₀)	hALR1 (IC ₅₀)	hALR1 / hALR2
1	8 nM	13,000 nM	1,200
2	10nM	11,000nM	1,100
3	5 nM	27,000 nM	5,400
4	8 nM	34,000 nM	4,250
5	6 nM	21,000 nM	3,500
6	8 nM	2,700 nM	340
7	12 nM	4,800 nM	400
8	7 nM	7,500 nM	1,100
9	11 nM	21,000 nM	1,900
10	5nM	13,000 nM	2,600
11	99 nM	5,600 nM	57

12	102 nM	10,000 nM	98
13	73 nM	13,000 nM	178
14	101 nM	16,000	160
15	53 nM	10,000	190
16	25 nM	6,200 nM	248
17	8 nM	41,000 nM	5,100
18	15 nM	>100 μ M	>6,700
19	30 nM	11,000 nM	370
20	7 nM	7,000 nM	1,000
21	14 nM	18,000 nM	1,300
22	9.1 nM	19,000 nM	2,100
23	9 nM	6,500 nM	720
24	1,040 nM	4,500 nM	4
25	160 nM	6,500 nM	41
26	17 nM	88,000 nM	5,200
27	52 nM	<5,000 nM	<96
28	5 nM	12,000 nM	2,400
29	11 nM	14,000	1,270
30	7.7 nM	21,000 nM	2,700
31	13 nM	9,700	746
32	660 nM	Not Tested	Not Tested
Tolrestat	13 nM	1,940 nM	149

The results show the superior potency, selectivity and efficacy of representative compounds of the invention. Such compounds are useful in the treatment of chronic complications arising from diabetes mellitus, such as, for example, diabetic cataracts, retinopathy and neuropathy. Accordingly, an aspect of the invention is treatment of such complications with the

inventive compounds; treatment includes both prevention and alleviation. The compounds are useful in the treatment of, for example, diabetic cataracts, retinopathy, nephropathy and neuropathy.

5 In a third, optional, set of experiments, the compounds can be assayed for their ability to normalize or reduce sorbitol accumulation in the sciatic nerve of streptozotocin-induced diabetic rats. The test methods employed to determine the efficacy are essentially those of Mylari, et al., *J. Med.*
10 *Chem.* 34: 108, 1991.

Example 34

The blood glucose lowering activity of the test compounds of this invention is demonstrated using the following
15 experiments with diabetic (db/db) mice.

The db/db (C57BL/KsJ) mouse exhibits many of the metabolic abnormalities that are associated with type 2 diabetes in humans. The mice are obese, extremely hyperglycemic and also hyperinsulinemic. Antihyperglycemic agents that are available
20 to man and also are effective in this model include metformin and troglitazone, both of which begin to demonstrate a beneficial effect in the db/db mice at doses above 100 mg/kg/day. Thus, compounds that are effective in this model are expected to be effective in humans.

Male db/db mice (8 weeks old) were obtained from Jackson Laboratories and were allowed to acclimate for 1 week before the experiment commenced. A sample of blood was collected from the tail after which plasma glucose was isolated by centrifugation and the glucose concentration was measured in the plasma enzymatically on the COBAS automated clinical analyzer equipped with a glucose kit that utilized hexokinase to quantitate the amount of glucose in a sample (Roche Diagnostic Systems, kit #47382). Mice with the lowest plasma glucose values were removed from the study and the remaining mice were randomized according to their individual plasma glucose values into 3 treatment groups (n=12 per group), control untreated db/db mice, 100 mg/kg/d compound treated db/db mice and 300 mg/kg/d compound treated db/db mice. The compound of Example 1 was administered in the diet by admixing the compound into the standard rodent powdered chow (Tekland LM-485 Mouse/Rat Sterilizable Diet 7012, Harlan Tekland).

Treatment with the compound was carried out for 4 weeks during which time blood glucose levels were measured weekly from the tail using the One Touch II blood glucose meter (Lifescan, Inc). The blood glucose values of mice in the compound treated groups was compared to the blood glucose values of mice from the control untreated group by an analysis of variance followed by Dunnett's Comparison Test (one-tailed).

The results in Table 1 show that the test compound of this invention lowers glucose in the diabetic db/db mouse over the 4 week study period. The mean percent change in glucose with drug treatment after four weeks of compound administration was 12% at a dose of 100 mg/kg/d and a 40% lowering of blood glucose at a dose of 300 mg/kg/d. This degree of glucose lowering is similar to what has been reported for troglitazone [+-5[[4-[(3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl) methoxy]phenyl)methyl]-2,4-thiazolidinedione] also known as CS-045 (Endocrinology, 1996, 137, 4189) and much better than that reported for metformin in this model. Thus the test compounds of this invention are well suited as antihyperglycemic agents.

Table 1: Blood glucose lowering

Group	Blood glucose (mg/dl)			
	Week 0	Week 2	Week 3	Week 4
Diabetic	299 ± 80	313 ± 90	305 ± 57	345 ± 46
Diabetic + test compound (100 mg/kg/d)	323 ± 79	284 ± 75	258 ± 60 ††	303 ± 79
Diabetic + test compound (300 mg/kg/d)	313 ± 52	240 ± 85	197 ± 77††	207 ± 77††

n=12 per group

†† p<0.05 compared to Diabetic

Data is given as mean + SD

Example 35

The assay described in this example is meant to determine whether the compounds of the instant invention would be effective in the treatment of elevated serum triglyceride levels in diabetic, as well as nondiabetic, patients. Tests are conducted to determine the effect of the compound of Example 1 on serum triglyceride levels in streptozotocin-induced diabetic rats. These animals represent a well-established diabetic model exhibiting most of the metabolic abnormalities associated with hyperglycemia, including hpertriglyceridemia, see Schnatz, et al., Diabetologia 8: 125, 1972.

Diabetes is induced in animals as follows: male Sprague-Dawley rats (150g), supplied by Harlan Teklad (Madison, WI), are allowed to acclimate for 1 week and water is supplied ad libitum. Food (7012CM, Harlan Teklad certified LM-485 mouse/rat) is removed at 1PM on the day prior to injection of streptozocin (STZ, Sigma cat no. S01230, lot no. 66H0468). STZ, 40 mg/kg, is prepared in 0.03 M citrate buffer, pH 4.5 and administered intraperitoneally after a 24-hr fast. Control animals receive citrate buffer.

Two hours after STZ injection, food is returned. Two days following STZ injection, blood glucose is measured and animals with <300 mg/dL are eliminated. Animals with blood glucose

levels ≥ 300 mg/dL are randomized into diabetic control and treated groups.

In all, three groups of animals are monitored and compared. The groups comprise a (nondiabetic) control group
5 (n=5); an untreated diabetic (control) group (n=7); and treated diabetic group (n=7). The daily dosages are administered at 10 AM by gavage as a single dose of the test compound in 2% Tween 90 in saline for 15 consecutive days. The nondiabetic and diabetic control groups are administered vehicle.

10 After the final dose (Day 8), all groups of animals are fasted for 4 hours and anesthetized with CO₂, and blood is collected by cardiac puncture into EDTA tubes. The plasma is separated from the red blood cells by centrifugation. Plasma triglyceride levels are quantitated on an automated COBAS
15 chemistry system utilizing the Roche reagent for Triglycerides (Cat #44119). This assay is a standard spectrophotometric enzyme assay that uses a Trinder reaction to measure the final product (Trinder, P., Ann Clin Biochem 6: 24-17, 1969). Statistical comparisons between groups employed a one-tailed t-
20 test.

Table 2 shows the results of the tests. As can be seen, administration of daily dosage of 10mg/kg significantly reduced the mean plasma triglyceride levels in treated animals 68% compared to the mean level for untreated diabetic animals. The
25 data clearly demonstrate the effectiveness of the test compound

in lowering serum triglyceride levels in diabetic animals; a property not generally associated with the ARI class. On the basis of these data, it is further to be expected that a similar effect would be produced in nondiabetic hosts with elevated triglyceride levels.

Table 2: Triglyceride lowering properties of test compound

Group	n	Plasma triglycerides (mg/dl)	Triglyceride lowering compared to diabetic
Control	5	62 ± 5	
Diabetic	7	335 ± 83*	
Diabetic + test compound (10 mg/kg/d)	7	149 ± 29†	68%

* p<0.01 compared to Control

†p<0.01 compared to the Diabetic

Data is given as mean + SEM

Example 36

The anti-angiogenic properties of the compounds of the invention are demonstrated in the following experiments using the rat aortic ring assay.

Rats (less than 6 weeks old, approx 150 grams) are individually sacrificed via carbon dioxide asphyxiation. The abdomen and thorax are opened along the midline with scissors using known sterile techniques. The animals are placed recumbent on their right side, to allow displacement of the

viscera. The abdominal and thoracic sections of the aorta are carefully separated from the dorsum by dissection along the longitudinal axis of the aorta. The isolated aorta is placed in a petri dish containing sterile, ice cold Hanks' balanced salt solution (Gibco BRL-Life Technologies, Rockville, MD) for further micro-dissection under a dissecting microscope. The luminal content of the aorta is dislodged by injection with Hanks, balanced salt solution via a syringe. Adherent adipose, loose connective tissue and segments of intercostal arteries are trimmed from the exterior of the aorta using sterile microsurgical instruments. The aorta is transferred to a clean petri dish containing fresh Hanks' balanced salt solution and the entire aorta is sectioned into 1 to 2 mm thick rings. The two end rings and any other rings which appear damaged are discarded. The aorta is maintained submerged in Hanks' balanced salt solution on ice while plating onto a 48-well plate.

Using a 48-well tissue culture plate, which is chilled on ice, in a tissue culture hood a 120 microliters of thawed Matrigel® (Becton Dickinson Labware, Bedford, MA) is plated onto each well using a sterile pipet tip. The Matrigel® is solidified by placing the culture plate for 30 minutes in a 37 °C humidified tissue culture incubator in the presence of 5% CO₂. A single aortic ring is placed on edge, with one of its two cut surfaces resting on the Matrigel®, at the center of

each well using a sterile curved forcep. The layout of the culture plate is such that the rings from multiple animals are placed in a single column on the plate. In this fashion, the experimental results represent observations based on six
5 animals (n=6). The aortic rings are completely embedded in Matrigel® by pipetting an additional 50 microliters of chilled Matrigel® over each ring, being careful not to disrupt proper ring orientation. The plate with the aortic rings is placed in an incubator at 37°C with 97% humidity for 6 days.

10 Each 48-well tissue culture plate has the following template: six negative wells, six positive wells, with the remaining wells used to evaluate various concentrations of the test compound in replicates of six. The six negative controls consisting of 1584 µL of human endothelial serum free media
15 (SFM) basal growth medium (Gibco-BRL-Life Technologies) and 16 µL of 100% sterile filtered DMSO. The six positive controls consist of 1484 µL serum free media (SFM) and 100 µL of endothelial cell growth supplement (ECGS, at a working concentration of 200 micrograms/ml) (Becton Dickinson Labware,
20 Bedford, MA), and 16 µL of 100% sterile filtered DMSO. Six wells, for each concentration of the test compound, consist of: 1484 µl SFM, 100 µl ECGS, 16 µl of test compounds dissolved in 100 % sterile filtered DMSO. Final concentration of DMSO in all wells is 1%. All test compounds are diluted 1:100 from their
25 stock concentrations.

Anti-angiogenic activity is verified using a double-cross over experimental design. The negative control (negative), the positive control (positive) and the experimental group labeled (prevention) each receive media changes every 24 hours for six
5 days. The content of the media changes are as previously described for each experimental group. The prevention group receives 50 micromolar concentration test compound for six consecutive days. The experimental group designated as Removal receives 50 micromolar concentration of test compound during
10 day 1, 2 and 3 after which the compound is removed by multiple rinses with fresh media. The aortic rings in the Removal group are cultured for an additional three days in the absence of the compound and treated identically to the positive control group on days 4, 5 and 6. The experimental group, labeled
15 Intervention, receives treatment identical to the positive control group for 1, 2 and 3 days and is then exposed to 50 microliters of test compound only on days 4, 5 and 6 in a fashion identical to the treatment received by the prevention group.

20 For quantitation of the angiogenic response, an inverted microscope (Zeiss, Axiovert 25) set at low illumination with full closure of the iris diaphragm to maximize depth-of-field is used. The microscope is coupled to a CCD camera (Cohu Inc.) for digital capture with a computer and each well of the 48-
25 well plate containing an aortic ring is digitally documented

At the completion of the study, the media is aspirated and Diff-Quik fixative (Dade-Behring) is added to each well as per manufacturer's instructions to preserve the specimens which are stored sealed and refrigerated at 4°C.

Table 3

Negative Control	0.72 ± 0.39 , n=6 **
Positive Control	2.95 ± 0.52 , n=6
Prevention (Days 1-6)	1.46 ± 0.48 , n=6 *
Removal (Days 4-6)	2.31 ± 0.71 , n=5
Intervention (Days 4-6)	1.97 ± 0.66 , n=5

20

Example 37


STZ treated diabetic minipigs having various wounds are administered the compound of Example 1. These animals are compared with control STZ diabetic minipigs also having wounds
5 but that are not treated with the compound. The animals administered the compound of Example 1 demonstrate a significant increase in the degree of wound healing. Accordingly, the compounds of the invention are capable of promoting wound healing in diabetic mammals.

10

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood
15 that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as
20 invention, the following claims conclude this specification.

What is claimed is:

1. A method for reducing elevated serum triglyceride levels, which method comprises administering to a mammal in need of such treatment an effective amount of a compound of the
5 formula:



wherein

A is a C₁-C₄ alkylene group optionally substituted with C₁-C₂ alkyl or mono- or disubstituted with halogen;

10 Z is a bond, O, S, C(O)NH, or C₁-C₃ alkylene optionally substituted with C₁-C₂ alkyl;


R₁ is hydrogen, alkyl having 1-6 carbon atoms, halogen, 2-, 3-, or 4-pyridyl, or phenyl, where the phenyl or pyridyl is optionally substituted with up to three groups selected
15 from halogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl, nitro, amino, or mono- or di(C₁-C₆)alkylamino;

R₂, R₃, R₄ and R₅ are each independently

hydrogen, halogen, nitro, or an alkyl group of 1-6 carbon atoms (which may be substituted with one or more
20 halogens);

OR₇, SR₇, S(O)R₇, S(O)₂N(R₇)₂, C(O)N(R₇)₂, or N(R₇)₂, wherein each R₇ is independently hydrogen, an alkyl group of 1-

6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-, 3-, or 4-pyridyl, each of which phenyl or heteroaryl is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenoxy where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; or a group of the formula



where

J is a bond, CH₂, oxygen, or nitrogen; and

each r is independently 2 or 3;

20 R₆ is hydroxy or a prodrug group;

R_a is hydrogen, C₁-C₆ alkyl, fluoro, or trifluoromethyl;

and Ar represents

a phenyl group optionally substituted with up to 5 groups
independently selected from halogen, an alkyl group of 1-6

carbon atoms (which may be substituted with one or more halogens), nitro, OR₁, SR₁, S(O)R₁, S(O)₂R₁, or N(R₁)₂, wherein R₁ is hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino, or the phenyl group may be condensed with benzo where the benzo is optionally substituted with one or two of halogen, cyano, nitro, trifluoromethyl, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl;

a heterocyclic 5-membered ring having one nitrogen, oxygen or sulfur, two nitrogens one of which may be replaced by oxygen or sulfur, or three nitrogens one of which may be replaced by oxygen or sulfur, said heterocyclic 5-membered ring substituted by one or two fluoro, chloro, (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo, cyano, nitro, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, one or two of fluoro, chloro, bromo, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio,

trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl or trifluoromethyl, or two fluoro or two trifluoromethyl with one hydroxy or one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one trifluoromethyl, or three fluoro, said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, bromo, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

a heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur, said heterocyclic 6-membered ring substituted by one or two (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo or trifluoromethylthio, or one or two of fluoro, chloro, bromo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, or trifluoromethyl, and said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;



said benzo-condensed heterocyclic 5-membered or 6-membered rings optionally substituted in the heterocyclic 5-membered or 6-membered ring by one of fluoro, chloro, bromo, methoxy, or trifluoromethyl;

oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms, with thiophene or

with furane, each optionally substituted by one of fluoro, chloro, bromo, trifluoromethyl, methylthio or methylsulfinyl;
imidazolopyridine or triazolopyridine optionally substituted by
5 one of trifluoromethyl, trifluoromethylthio, bromo, or
(C₁-C₆)alkoxy, or two of fluoro or chloro;
thienothiophene or thienofuran optionally substituted by one of
fluoro, chloro or trifluoromethyl;
thienotriazole optionally substituted by one of chloro or
10 trifluoromethyl;
naphthothiazole; naphthoxazole; or thienoisothiazole.

2. A method according to claim 1, wherein Ar is aryl or heteroaryl, each of which is substituted with up to four groups
15 independently selected from hydrogen, fluorine, chlorine, bromine, trifluoromethyl and nitro.

3. A method according to claim 1, wherein Ar is a substituted phenyl of Formula II or a substituted benzothiazole
20 of Formula III



wherein R_8 , R_8' , R_9 , R_9' , R_{10} , R_{11} , R_{12} , R_{13} and R_{14} are independently hydrogen, fluorine, chlorine, bromine, trifluoromethyl or nitro.

5 4. A method according to claim 3, wherein A is methylene and Z is a bond.

 5. A method according to claim 3, wherein R_8 is hydrogen and Z is a bond.

10

 6. A method according to claim 3, wherein A is methylene, R_8 is hydrogen, and Z is a bond.

 7. A method according to claim 6, wherein Ar is a
15 substituted benzothiazole of Formula III.

 8. A method according to claim 7, wherein at least one of R_{11} , R_{12} , R_{13} , and R_{14} is trifluoromethyl.

20 9. A method according to claim 8, wherein R_{12} is trifluoromethyl.

 10. A method according to claim 7, wherein R_{11} , R_{12} , and R_{14} are fluorines and R_{13} is hydrogen.

25

11. A method according to claim 10, wherein R_6 is hydrogen.

12. A method according to claim 10, wherein R_6 is C_1-C_6
5 alkyl.

13. A method according to claim 6, wherein Ar is a substituted phenyl of Formula II.

10 14. A method according to claim 13, wherein at least one of R_8 , R_8 , R_9 , R_9 , R_{10} is trifluoromethyl.

15 15. A method according to claim 14, wherein R_9 is trifluoromethyl.

16 16. A method according to claim 15, wherein R_8 , R_8 , R_9 , R_9 , R_{10} are fluorines and R_{11} is hydrogen.

20 17. A method according to claim 16, wherein R_6 is hydrogen.

25 18. A method according to claim 16, wherein R_6 is C_1-C_6 alkyl.

19. A method according to claim 1, which is 3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid, ethyl Ester.

5 20. A method according to claim 1, which is 3-(4,5,7-trifluorobenzothiazol-2-yl) methyl-indole-*N*-acetic acid.

21. A method according to claim 1, which is 5-chloro-3-(4,5,7-Trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.
10

22. A method according to claim 1, which is 5-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

23. A method according to claim 1, which is 2-methyl-3-(4,5,7 trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.
15

24. A method according to claim 1, which is 5-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

20 25. A method according to claim 1, which is 7-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

26. A method according to claim 1, which is 6-chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.
25

27. A method according to claim 1, which is 5-benzyloxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

5 28. A method according to claim 1, which is 6-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

29. A method according to claim 1, which is 5-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

10

30. A method according to claim 1, which is 6-methyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

31. A method according to claim 1, which is 3-(5-
15 trifluoromethylbenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

32. A method according to claim 1, which is 5-Methyl-3-(5-Trifluoromethylbenzothiazol-2-yl)methyl-indole-*N*-acetic acid.

20

33. A method according to claim 1, which is 3-(3-nitrophenyl)methyl-indole-*N*-acetic acid.

34. A method according to claim 1, which is 3-(3-
25 nitrophenyl)methyl-indole-*N*-acetic acid, ethyl ester.

35. A method according to claim 1, which is 3-(3-nitrophenyl)methyl-indole-N-acetic acid.

5 36. A method according to claim 1, which is 2-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

37. A method according to claim 1, which is 5-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

10

38. A method according to claim 1, which is 6-phenyl-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

39. A method according to claim 1, which is 5-morpholino-
15 3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic
acid.

40. A method according to claim 1, which is 6-morpholino-
3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic
20 acid.

41. A method according to claim 1, which is 5-phenoxy-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

42. A method according to claim 1, which is 7-fluoro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

43. A method according to claim 1, which is 7-bromo-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

44. A method according to claim 1, which is -chloro-3-(4,5,7-trifluorobenzothiazol-2-yl)methyl-indole-N-acetic acid.

45. A method according to claim 1, which is 3-[[5-Fluorobenzothiazole-2-yl)methyl]-indole-N-acetic acid.

46. A method according to claim 1, which is 3-[[6-Fluorobenzothiazole-2-yl)methyl]-indole-N-acetic acid.

15


47. A method according to claim 3, wherein Ar is a substituted benzothiazole of Formula III, R_{12} is trifluoromethyl, A is methylene, methylene substituted with a methyl group, or ethylene, and R_2 , R_3 , R_4 and R_5 , in combination, represent one of bromo, cyano or nitro, one or two of fluoro, chloro, hydroxy, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, or trifluoromethyl, or two fluoro or two methyl with one hydroxy or one (C_1-C_6) alkoxy, or one or, preferably, two fluoro and one methyl, or three fluoro groups.

48. A method according to claim 9, wherein the mammal is diabetic.

49. A method according to claim 9, wherein the mammal is
5 non-diabetic.

50. A method according to claim 1, wherein the compound
is administered in a combination therapy along with at least
one second agent used to treat hyperglycemia, hyperlipidemia,
10 or diabetic complications.

51. A method for reducing elevated serum glucose levels,
which method comprises administering to a mammal in need of
such treatment an effective amount of a compound of the
15 formula:



wherein

A is a C₁-C₄ alkylene group optionally substituted with C₁-C₂
alkyl or mono- or disubstituted with halogen;

20 Z is a bond, O, S, C(O)NH, or C₁-C₃ alkylene optionally
substituted with C₁-C₂ alkyl;

R₁ is hydrogen, alkyl having 1-6 carbon atoms, halogen, 2-, 3-,
or 4-pyridyl, or phenyl, where the phenyl or pyridyl is
optionally substituted with up to three groups selected
from halogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl, nitro,
5 amino, or mono- or di(C₁-C₆)alkylamino;

R₂, R₃, R₄ and R₅ are each independently


hydrogen, halogen, nitro, or an alkyl group of 1-6 carbon
atoms (which may be substituted with one or more
halogens);

10 OR₇, SR₇, S(O)R₇, S(O)₂N(R₇)₂, C(O)N(R₇)₂, or N(R₇)₂, wherein
each R₇ is independently hydrogen, an alkyl group of 1-
6 carbon atoms (which may be substituted with one or
more halogens) or benzyl, where the phenyl portion is
optionally substituted with up to three groups
15 independently selected from halogen, C₁-C₆ alkyl, C₁-C₆
alkoxy, amino, and mono- or di(C₁-C₆)alkylamino;

phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-,
3-, or 4-pyridyl, each of which phenyl or heteroaryl
is optionally substituted with up to three groups
independently selected from halogen, C₁-C₆ alkyl, C₁-C₆
20 alkoxy, amino, and mono- or di(C₁-C₆)alkylamino;

phenoxy where the phenyl portion is optionally substituted
with up to three groups independently selected from
halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or
25 di(C₁-C₆)alkylamino; or

a group of the formula



where

J is a bond, CH₂, oxygen, or nitrogen; and

5 each r is independently 2 or 3;

R₆ is hydroxy or a prodrug group;

R_a is hydrogen, C₁-C₆ alkyl, fluoro, or trifluoromethyl;

and Ar represents

a phenyl group optionally substituted with up to 5 groups

10 independently selected from halogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens), nitro, OR₇, SR₇, S(O)R₇, S(O)₂R₇ or N(R₇)₂, wherein

R₇ is hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl,

15 where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino, or the phenyl group may be condensed with

benzo where the benzo is optionally substituted with one or two of halogen, cyano, nitro, trifluoromethyl, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio,

20 trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl;

a heterocyclic 5-membered ring having one nitrogen, oxygen or sulfur, two nitrogens one of which may be replaced by oxygen or sulfur, or three nitrogens one of which may be replaced by oxygen or sulfur, said heterocyclic 5-membered ring substituted by one or two fluoro, chloro, (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo, cyano, nitro, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, one or two of fluoro, chloro, bromo, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl or trifluoromethyl, or two fluoro or two trifluoromethyl with one hydroxy or one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one trifluoromethyl, or three fluoro, said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, bromo, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

a heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur, said heterocyclic 6-membered ring substituted by one or two (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo or trifluoromethylthio, or one or two of fluoro, chloro,

bromo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, or trifluoromethyl, and said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

said benzo-condensed heterocyclic 5-membered or 6-membered rings optionally substituted in the heterocyclic 5-membered or 6-membered ring by one of fluoro, chloro, bromo, methoxy, or trifluoromethyl;

oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms, with thiophene or with furane, each optionally substituted by one of fluoro, chloro, bromo, trifluoromethyl, methylthio or methylsulfinyl;


imidazolopyridine or triazolopyridine optionally substituted by one of trifluoromethyl, trifluoromethylthio, bromo, or (C₁-C₆)alkoxy, or two of fluoro or chloro;

thienothiophene or thienofuran optionally substituted by one of fluoro, chloro or trifluoromethyl;

thienotriazole optionally substituted by one of chloro or trifluoromethyl;

naphthothiazole; naphthoxazole; or thienoisothiazole.

52. A method for inhibiting angiogenesis, which method comprises administering to a mammal in need of such treatment an effective amount of a compound of the formula:



5 wherein

A is a C₁-C₄ alkylene group optionally substituted with C₁-C₂ alkyl or mono- or disubstituted with halogen;


Z is a bond, O, S, C(O)NH, or C₁-C₃ alkylene optionally substituted with C₁-C₂ alkyl;

10 R₁ is hydrogen, alkyl having 1-6 carbon atoms, halogen, 2-, 3-, or 4-pyridyl, or phenyl, where the phenyl or pyridyl is optionally substituted with up to three groups selected from halogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl, nitro, amino, or mono- or di(C₁-C₆)alkylamino;

15 R₂, R₃, R₄ and R₅ are each independently hydrogen, halogen, nitro, or an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens);

OR₇, SR₇, S(O)R₇, S(O)₂N(R₇)₂, C(O)N(R₇)₂, or N(R₇)₂, wherein
20 each R₇ is independently hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is

optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-, 3-, or 4-pyridyl, each of which phenyl or heteroaryl is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenoxy where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; or a group of the formula



where J is a bond, CH₂, oxygen, or nitrogen; and each r is independently 2 or 3; R₆ is hydroxy or a prodrug group; R_a is hydrogen, C₁-C₆ alkyl, fluoro, or trifluoromethyl; and Ar represents a phenyl group optionally substituted with up to 5 groups independently selected from halogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens), nitro, OR₇, SR₇, S(O)R₇, S(O)₂R₇, or N(R₇)₂, wherein

R₇ is hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino, or the phenyl group may be condensed with benzo where the benzo is optionally substituted with one or two of halogen, cyano, nitro, trifluoromethyl, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl;

a heterocyclic 5-membered ring having one nitrogen, oxygen or sulfur, two nitrogens one of which may be replaced by oxygen or sulfur, or three nitrogens one of which may be replaced by oxygen or sulfur, said heterocyclic 5-membered ring substituted by one or two fluoro, chloro, (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo, cyano, nitro, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, one or two of fluoro, chloro, bromo, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl or trifluoromethyl,

or two fluoro or two trifluoromethyl with one hydroxy or one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one trifluoromethyl, or three fluoro, said pyridyl, furyl or thienyl optionally substituted in the 3-position by
5 fluoro, chloro, bromo, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

a heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur, said heterocyclic 6-membered ring substituted by one or two (C₁-C₆)alkyl or phenyl, or condensed with
10 benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo or trifluoromethylthio, or one or two of fluoro, chloro, bromo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, or trifluoromethyl,
15 and said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

said benzo-condensed heterocyclic 5-membered or 6-membered rings optionally substituted in the heterocyclic 5-
20 membered or 6-membered ring by one of fluoro, chloro, bromo, methoxy, or trifluoromethyl;

oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms, with thiophene or with furane, each optionally substituted by one of fluoro,

chloro, bromo, trifluoromethyl, methylthio or
methylsulfinyl;

imidazolopyridine or triazolopyridine optionally substituted by
one of trifluoromethyl, trifluoromethylthio, bromo, or


5 (C₁-C₆)alkoxy, or two of fluoro or chloro;

thienothiophene or thienofuran optionally substituted by one of
fluoro, chloro or trifluoromethyl;

thienotriazole optionally substituted by one of chloro or
trifluoromethyl;

10 naphthothiazole; naphthoxazole; or thienoisothiazole.

53. A method for reducing elevated serum glucose and
triglyceride levels, which method comprises administering to a
mammal in need of such treatment an effective amount of a
15 compound of the formula:



wherein

A is a C₁-C₄ alkylene group optionally substituted with C₁-C₂
alkyl or mono- or disubstituted with halogen;

20 Z is a bond, O, S, C(O)NH, or C₁-C₃ alkylene optionally
substituted with C₁-C₂ alkyl;

R_1 is hydrogen, alkyl having 1-6 carbon atoms, halogen, 2-, 3-,
or 4-pyridyl, or phenyl, where the phenyl or pyridyl is
optionally substituted with up to three groups selected
from halogen, hydroxy, C_1-C_6 alkoxy, C_1-C_6 alkyl, nitro,
5 amino, or mono- or di(C_1-C_6)alkylamino;

R_2 , R_3 , R_4 and R_5 are each independently


hydrogen, halogen, nitro, or an alkyl group of 1-6 carbon
atoms (which may be substituted with one or more
halogens);

10 OR_7 , SR_7 , $S(O)R_7$, $S(O)_2N(R_7)_2$, $C(O)N(R_7)_2$, or $N(R_7)_2$, wherein
each R_7 is independently hydrogen, an alkyl group of 1-
6 carbon atoms (which may be substituted with one or
more halogens) or benzyl, where the phenyl portion is
optionally substituted with up to three groups
15 independently selected from halogen, C_1-C_6 alkyl, C_1-C_6
alkoxy, amino, and mono- or di(C_1-C_6)alkylamino;

phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-,
3-, or 4-pyridyl, each of which phenyl or heteroaryl
is optionally substituted with up to three groups
20 independently selected from halogen, C_1-C_6 alkyl, C_1-C_6
alkoxy, amino, and mono- or di(C_1-C_6)alkylamino;

phenoxy where the phenyl portion is optionally substituted
with up to three groups independently selected from
halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, amino, and mono- or
25 di(C_1-C_6)alkylamino; or

a group of the formula



where

J is a bond, CH₂, oxygen, or nitrogen; and

5 each r is independently 2 or 3;

R₆ is hydroxy or a prodrug group;

R_a is hydrogen, C₁-C₆ alkyl, fluoro, or trifluoromethyl;

and Ar represents

a phenyl group optionally substituted with up to 5 groups

10 independently selected from halogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens), nitro, OR₇, SR₇, S(O)R₇, S(O)₂R₇, or N(R₇)₂, wherein

R₇ is hydrogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens) or benzyl,

15 where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino, or the phenyl group may be condensed with

20 benzo where the benzo is optionally substituted with one or two of halogen, cyano, nitro, trifluoromethyl, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl;

a heterocyclic 5-membered ring having one nitrogen, oxygen or sulfur, two nitrogens one of which may be replaced by oxygen or sulfur, or three nitrogens one of which may be replaced by oxygen or sulfur, said heterocyclic 5-membered ring substituted by one or two fluoro, chloro, (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo, cyano, nitro, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, one or two of fluoro, chloro, bromo, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl or trifluoromethyl, or two fluoro or two trifluoromethyl with one hydroxy or one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one trifluoromethyl, or three fluoro, said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, bromo, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

a heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur, said heterocyclic 6-membered ring substituted by one or two (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo or trifluoromethylthio, or one or two of fluoro, chloro,

bromo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, or trifluoromethyl, and said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

said benzo-condensed heterocyclic 5-membered or 6-membered rings optionally substituted in the heterocyclic 5-membered or 6-membered ring by one of fluoro, chloro, bromo, methoxy, or trifluoromethyl;

oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms, with thiophene or with furane, each optionally substituted by one of fluoro, chloro, bromo, trifluoromethyl, methylthio or methylsulfinyl;

imidazolopyridine or triazolopyridine optionally substituted by one of trifluoromethyl, trifluoromethylthio, bromo, or (C₁-C₆)alkoxy, or two of fluoro or chloro;


thienothiophene or thienofuran optionally substituted by one of fluoro, chloro or trifluoromethyl;

thienotriazole optionally substituted by one of chloro or trifluoromethyl;

naphthothiazole; naphthoxazole; or thienoisothiazole.

54. A method for promoting wound healing in mammals, which method comprises administering to a mammal in need of


such treatment an effective amount of a compound of the formula:



wherein

- 5 A is a C₁-C₄ alkylene group optionally substituted with C₁-C₂ alkyl or mono- or disubstituted with halogen;
- Z is a bond, O, S, C(O)NH, or C₁-C₃ alkylene optionally substituted with C₁-C₂ alkyl;
- 10 R₁ is hydrogen, alkyl having 1-6 carbon atoms, halogen, 2-, 3-, or 4-pyridyl, or phenyl, where the phenyl or pyridyl is optionally substituted with up to three groups selected from halogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl, nitro, amino, or mono- or di(C₁-C₆)alkylamino;
- R₂, R₃, R₄ and R₅ are each independently
- 15 hydrogen, halogen, nitro, or an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens);
- OR₇, SR₇, S(O)R₇, S(O)₂N(R₇)₂, C(O)N(R₇)₂, or N(R₇)₂, wherein each R₇ is independently hydrogen, an alkyl group of 1-
- 20 6 carbon atoms (which may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups

independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenyl or heteroaryl such as 2-, 3- or 4-imidazolyl or 2-, 3-, or 4-pyridyl, each of which phenyl or heteroaryl is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; phenoxy where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino; or a group of the formula



where

J is a bond, CH₂, oxygen, or nitrogen; and each r is independently 2 or 3;

R₆ is hydroxy or a prodrug group;

R₈ is hydrogen, C₁-C₆ alkyl, fluoro, or trifluoromethyl;

and Ar represents

a phenyl group optionally substituted with up to 5 groups independently selected from halogen, an alkyl group of 1-6 carbon atoms (which may be substituted with one or more halogens), nitro, OR₇, SR₇, S(O)R₇, S(O)₂R₇ or N(R₇)₂ wherein R₇ is hydrogen, an alkyl group of 1-6 carbon atoms (which

may be substituted with one or more halogens) or benzyl, where the phenyl portion is optionally substituted with up to three groups independently selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, amino, and mono- or di(C₁-C₆)alkylamino, or the phenyl group may be condensed with benzo where the benzo is optionally substituted with one or two of halogen, cyano, nitro, trifluoromethyl, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl;

a heterocyclic 5-membered ring having one nitrogen, oxygen or sulfur, two nitrogens one of which may be replaced by oxygen or sulfur, or three nitrogens one of which may be replaced by oxygen or sulfur, said heterocyclic 5-membered ring substituted by one or two fluoro, chloro, (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl, said phenyl or benzo optionally substituted by one of iodo, cyano, nitro, perfluoroethyl, trifluoroacetyl, or (C₁-C₆)alkanoyl, one or two of fluoro, chloro, bromo, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, trifluoromethoxy, trifluoromethylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl or trifluoromethyl, or two fluoro or two trifluoromethyl with one hydroxy or

one (C₁-C₆)alkoxy, or one or, preferably, two fluoro and one trifluoromethyl, or three fluoro, said pyridyl, furyl or thienyl optionally substituted in the 3-position by fluoro, chloro, bromo, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

5 a heterocyclic 6-membered ring having one to three nitrogen atoms, or one or two nitrogen atoms and one oxygen or sulfur, said heterocyclic 6-membered ring substituted by one or two (C₁-C₆)alkyl or phenyl, or condensed with benzo, or substituted by one of pyridyl, furyl or thienyl,
10 said phenyl or benzo optionally substituted by one of iodo or trifluoromethylthio, or one or two of fluoro, chloro, bromo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, or trifluoromethyl, and said pyridyl, furyl or thienyl optionally substituted
15 in the 3-position by fluoro, chloro, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

said benzo-condensed heterocyclic 5-membered or 6-membered rings optionally substituted in the heterocyclic 5-membered or 6-membered ring by one of fluoro, chloro,
20 bromo, methoxy, or trifluoromethyl;

oxazole or thiazole condensed with a 6-membered aromatic group containing one or two nitrogen atoms, with thiophene or with furane, each optionally substituted by one of fluoro, chloro, bromo, trifluoromethyl, methylthio or
25 methylsulfinyl;

imidazolopyridine or triazolopyridine optionally substituted by
one of trifluoromethyl, trifluoromethylthio, bromo, or
(C₁-C₆)alkoxy, or two of fluoro or chloro;
thienothiophene or thienofuran optionally substituted by one of
5 fluoro, chloro or trifluoromethyl;
thienotriazole optionally substituted by one of chloro or
trifluoromethyl;
naphthothiazole; naphthoxazole; or thienoisothiazole.

10 55. A method according to claim 54, wherein the mammal is
diabetic.

56. A method according to claim 55, wherein the mammal is
human.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/404, 31/405, 31/428, A61P 3/06, 3/10, 9/14, 17/02, // A61K 31/00, 31:404, 31:405, 31:428	A3	(11) International Publication Number: WO 00/32180 (43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/US99/28483 (22) International Filing Date: 1 December 1999 (01.12.99) (30) Priority Data: 60/110,395 1 December 1998 (01.12.98) US (71) Applicant (for all designated States except US): THE INSTI- TUTES FOR PHARMACEUTICAL DISCOVERY, INC. [US/US]; 23 Business Park Drive, Branford, CT 06405 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SREDY, Janet [US/US]; 98 Trumbull Avenue, Milford, CN 06460 (US). JACOT, Jorge [US/US]; 2 Katie-Joe Lane, Branford, CN 06405 (US). (74) Agent: SARUSSI, Steven, J.; McDonnell, Bochnen, Hulbert & Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 16 November 2000 (16.11.00)
(54) Title: ANTIHYPERTRIGLYCERIDEMIC, ANTIHYPERGLYCEMIC, ANTI-ANGIOGENIC AND WOUND HEALING SUBSTITUTED INDOLEALKANOIC ACIDS (57) Abstract Disclosed are methods of reducing serum glucose and triglyceride levels and for inhibiting angiogenesis, the methods comprising administration of substituted indolealkanoic acids to patients in need of such treatment. Also disclosed are such compounds useful in the treatment of angiogenesis, hyperglycemia, hyperlipidemia and chronic complications arising from diabetes mellitus. Also disclosed are pharmaceutical compositions containing the compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/28483

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/404 A61K31/405 A61K31/428 A61P3/06 A61P3/10
A61P9/14 A61P17/02 //A61K31/00,31:404,31:405,31:428

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 50268 A (THE INSTITUTES OF PHARMACEUTICAL DISCOVERY) 7 October 1999 (1999-10-07) the whole document ---	1-56
A	WO 96 26207 A (NISSAN CHEMICAL INDUSTRIES) 29 August 1996 (1996-08-29) see the whole document, especially page 4, page 170 lines 1-19, "test example 1" on pages 225-226, "test example 2" on pages 228-229; page 232 ---	1-56
A	EP 0 624 369 A (PFIZER) 17 November 1994 (1994-11-17) the whole document ---	1-18
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 August 2000

Date of mailing of the international search report

07/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gac, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/28483

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 0, no. 0 & JP 09 165371 A (SANKYO CO LTD), 24 June 1997 (1997-06-24) abstract -----	1-6, 13-18, 51,53-56
A	US 4 960 785 A (H. R. HOWARD ET AL.) 2 October 1990 (1990-10-02) column 1 -column 2 see formula I column 7, line 35 - line 50 -----	51,53-56
A	EP 0 054 417 A (PFIZER CORPORATION) 23 June 1982 (1982-06-23) page 2, line 8 - line 9 page 10, line 14 - line 24 page 11, line 23 - line 26 page 12, line 1 - line 12 -----	52,54-56
A	EP 0 705 607 A (TAKEDA CHEM. IND.) 10 April 1996 (1996-04-10) the whole document -----	1,50,53

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Present claims 1-18,47-56 relate to an extremely large number of possible compounds/methods. In fact, the claims contain so many options, variables or possible permutations that a lack of clarity (and/or conciseness) within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search on the whole scope of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear and supported, namely on the specific compounds mentioned in claims 19-46 and to compounds of examples 26-32 with due regard to the general idea underlying the invention.

The expression ... "along with at least one second agent used to treat hyperglycemia, hyperlipidemia or diabetic complications" in claim 50 is neither clear nor supported to provide for a definition of structural compounds and was therefore not searched.

Please note that claim 35 is identical to claim 33 and should therefore be deleted.

In claim 44, correction of the typing error ("-chloro-3-(.....)" should be : " 7-chloro-3-(.....)") will be necessary.

Claims searched completely : claims 19-46

Claims searched incompletely : claims 1-18, 47-56

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/28483

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9950268	A	07-10-1999	AU	3459599 A	18-10-1999
WO 9626207	A	29-08-1996	AU	4731196 A	11-09-1996
			JP	9235284 A	09-09-1997
			ZA	9601478 A	28-08-1996
EP 624369	A	17-11-1994	US	5391551 A	21-02-1995
			AU	657444 B	09-03-1995
			AU	6196694 A	01-12-1994
			CA	2123059 A,C	11-11-1994
			CN	1103783 A,B	21-06-1995
			IL	109517 A	16-08-1998
			JP	8165241 A	25-06-1996
			KR	140146 B	01-06-1998
			ZA	9403203 A	09-11-1995
JP 09165371	A	24-06-1997	NONE		
US 4960785	A	02-10-1990	EP	0252713 A	13-01-1988
			US	5064852 A	12-11-1991
EP 54417	A	23-06-1982	AR	229110 A	15-06-1983
			AU	525296 B	28-10-1982
			AU	7848881 A	24-06-1982
			CA	1143737 A	29-03-1983
			CS	228527 B	14-05-1984
			DD	202290 A	07-09-1983
			DE	3167879 D	31-01-1985
			DK	552581 A	16-06-1982
			ES	507960 D	01-10-1983
			ES	8308872 A	16-12-1983
			FI	814003 A	16-06-1982
			GR	76355 A	06-08-1984
			IE	51949 B	29-04-1987
			JP	57181082 A	08-11-1982
			NO	814264 A	16-06-1982
			PH	17576 A	01-10-1984
			PL	234253 A	23-05-1983
			PT	74123 A,B	01-01-1982
			US	4363912 A	14-12-1982
			YU	293481 A	31-10-1984
			ZA	8108665 A	27-10-1982
EP 705607	A	10-04-1996	AU	703422 B	25-03-1999
			AU	3308195 A	18-04-1996
			CA	2160092 A	08-04-1996
			CN	1128657 A	14-08-1996
			FI	954773 A	08-04-1996
			HU	73422 A	29-07-1996
			JP	8157369 A	18-06-1996
			NO	953961 A	09-04-1996
			NZ	280165 A	27-07-1997
			US	5726306 A	10-03-1998
			US	5885979 A	23-03-1999

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.